

B4

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
20 September 2001 (20.09.2001)

PCT

(10) International Publication Number
WO 01/68600 A2

BEST AVAILABLE COPY

- (51) International Patent Classification⁷: C07D 207/00 SHEN, Yaping [CA/CA]; 21-2458 Pitt River Road, Port Coquitlam, British Columbia V3C 6J7 (CA).
- (21) International Application Number: PCT/CA01/00345
- (22) International Filing Date: 16 March 2001 (16.03.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/190,337 16 March 2000 (16.03.2000) US
- (71) Applicant (for all designated States except US): INFLAZYME PHARMACEUTICALS LIMITED [CA/CA]; Suite 425, 5600 Parkwood Way, Richmond, British Columbia V6V 2M2 (CA).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): LAUENER, Ronald, W. [CA/CA]; 818 Massey Street, New Westminster, British Columbia V3L 4S9 (CA). BURGOYNE, David, L. [CA/CA]; 4715 53A Street, Delta, British Columbia V4K 3V4 (CA). REBSTEIN, Patrick, J. [CA/CA]; 2659 West 14th Avenue, Vancouver, British Columbia V6K 2W8 (CA). MACKENZIE, Lloyd, F. [CA/CA]; 10864 129th Street, Surrey, British Columbia V3T 3N8 (CA). ZHOU, Yuanlin [CA/CA]; 11253 Daniels Road, Richmond, British Columbia V6X 1M5 (CA).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:
— without international search report and to be republished upon receipt of that report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



WO 01/68600 A2

(54) Title: BENZYLATED PDE4 INHIBITORS

(57) Abstract: The present invention is directed to a method for reducing the emetogenic effects of PDE inhibitors, and more particularly is directed to compounds having PDE4 inhibition activity with little or no emetogenic side-effects, and chemical methods including benzylation for preparing such compounds. A benzyl group may be attached to either a carbon or nitrogen atom of a PDE4 inhibitor. Suitable benzylation chemistry is to extract a hydrogen from a PDE4 inhibitor, preferably with a base, and then react the resulting nucleophilic PDE4 inhibitor with a benzylating agent, e.g. benzyl bromide or a derivative thereof.

BENZYLATED PDE4 INHIBITORS

TECHNICAL FIELD

The present invention is directed to a method for reducing the emetogenic effects of PDE inhibitors, and more particularly is directed to compounds having PDE4 inhibition activity with little or no emetogenic side-effects, and chemical methods including benzylation for preparing such compounds.

BACKGROUND OF THE INVENTION

Enzymes known as phosphodiesterases (PDEs) function *in vivo* to hydrolytically cleave the 3'-phosphodiester bond of cyclic nucleotides to thereby form the corresponding 5'-monophosphate. For instance, certain PDEs can hydrolyze the 3'-phosphodiester bond of adenosine 3',5'-cyclic monophosphate (cAMP) so as to form 5'-adenosine monophosphate (5'-AMP), and/or can hydrolyze the 3'-phosphodiester bond of guanosine 3',5'-cyclic monophosphate (cGMP) so as to form 5'-guanosine monophosphate (5'-GMP). These cyclic nucleotides exert a significant impact on cellular processes by, for example, converting inactive protein kinase enzymes into an active form. The active form of the protein kinase catalyzes various phosphorylation processes that impact on fundamental cellular processes including transcriptional regulation, ion channel function, and signaling protein activity.

Researchers investigating PDEs generally agree that there are at least eleven distinct PDE families, differentiated on the basis of amino acid sequence, substrate specificity and sensitivity to endogenous and exogenous regulators. These families are commonly known as PDE1 through PDE11. In addition, researchers found that cyclic nucleotide concentration is a significant factor in the course of the *in vivo* inflammatory response. Accordingly, much research has been directed to methods for influencing the concentration of cyclic nucleotides as a means to influence the inflammatory response, and particular attention has been directed at PDE4 activity. One promising area of research is the development of small organic molecules that inhibit PDE activity. By inhibiting PDE activity, these small molecules reduce the amount of cyclic nucleotide that is converted into the (inactive) corresponding 5'-monophosphate, thereby elevating cyclic nucleotide concentration, and indirectly increasing protein kinase activity within the cell.

Many major pharmaceutical companies are working to develop specific small organic molecules into pharmaceutical compositions that function as PDE inhibitors. ROLIPRAM™ (Schering AG) is an example of an early attempt to develop

This Page Blank (uspto)

such a composition directed to PDE4. However, while ROLIPRAM™ exhibited marked anti-inflammatory activity, it was also found to demonstrate unwanted side effects including emesis (also known as nausea and vomiting) and potentiation of gastric acid secretion. These undesired side effects caused ROLIPRAM™ to be
5 withdrawn from development as an anti-inflammatory pharmaceutical. An understanding of the cause of these side-effects, and approaches to mitigate them, subsequently became topics of intense study.

It is now recognized that PDE4 exists in two distinct forms, *i.e.*, two conformers. One conformer, known variously as HPDE4 or HARB, is particularly
10 prevalent in the gastrointestinal tract and central nervous system, has a high affinity for ROLIPRAM™ (*i.e.*, has a High Affinity ROLIPRAM™ Binding Site, "HARBS"), and is considered responsible for the unwanted side-effects. The other conformer is known variously as LPDE4 or LARB, is found in immunocompetent cells, and has a low affinity for ROLIPRAM™. Researchers are seeking to develop small molecules that inhibit the
15 catalytic activity of LPDE4 rather than bind to HPDE4, *i.e.*, molecules that have a low LPDE4:HPDE4 ratio where the numerator and denominator are the appropriate IC₅₀ values. In other words, researchers are seeking so-called "second generation" inhibitor molecules that interact with the LPDE4 catalytic site of PDE4, rather than the HPDE4 ROLIPRAM™ binding site, to provide desirable anti-inflammatory effect without
20 unwanted side-effects such as emesis.


The present invention is directed to fulfilling this need in the art, and providing further related advantages as set forth more completely herein.

For additional and more detailed discussion of PDE enzymes, including the history of their discovery, their characterization and classification, their *in vivo*
25 activity, their inhibition by small organic molecules, and current clinical efforts directed to providing pharmaceutical compositions containing these small molecules, *see, e.g.*, Burnouf, C. et al. "Phosphodiesterase 4 Inhibitors" *Annual Reports in Medicinal Chemistry*, Vol. 33, Chap. 10, pp 91-109, 1998 (Bristol, J.A., ed.); Essayan, D. M. "Cyclic Nucleotide Phosphodiesterase (PDE) Inhibitors and Immunomodulation"
30 *Biochemical Pharmacology* 57:965-973, 1999; Souness, J. E. and Foster, M. "Potential of phosphodiesterase type IV inhibitors in the treatment of rheumatoid arthritis" *Idrugs* 1(5):541-553, 1998; Souness, J.E. et al. "Immunosuppressive and anti-inflammatory effect of cAMP phosphodiesterase (PDE) type 4 inhibitors" *Immunopharmacology* 47: 127-162, 2000; and Torphy, T. J. "Phosphodiesterase Isozymes" *Am J. Respir. Crit.*
35 *Care Med.* 157:351-370, 1998, as well as the numerous references cited in these articles.

SUMMARY OF THE INVENTION

In one aspect, the present invention provides a method for improving the therapeutic ratio of PDE4 inhibitors through reduction or elimination of emetogenic properties while allowing retention of PDE activity. In one aspect, the method comprises benzylation of a PDE4 inhibitor, wherein the benzylation places a benzyl group on a carbon atom of a PDE4 inhibitor, to provide a benzylated PDE4 inhibitor. In one aspect, the PDE4 inhibitor is ROLIPRAM™, while in another aspect the PDE4 inhibitor is ARIFLO™. The benzylation may replace an extractable hydrogen of the PDE4 inhibitor with a benzyl group. For instance, the PDE4 inhibitor may be reacted with a base in order to extract the extractable hydrogen, followed by reaction with a benzylating agent. In one aspect, the PDE4 inhibitor has a carbonyl group and the extractable hydrogen is located alpha to (*i.e.*, next to) the carbonyl group. In another aspect, the PDE4 inhibitor has a carbonyl group and benzyl group is located beta to (*i.e.*, with one intervening atom between) the carbonyl group. In another aspect, the PDE4 inhibitor has a carbonyl group and benzyl group is located gamma to (*i.e.*, with two intervening atoms between) the carbonyl group. The extractable hydrogen may be removed under chemical conditions and replaced with a benzyl group.

In one aspect, the benzylated PDE4 inhibitor comprises at least one benzyl group (Bzl) of the formula




wherein each of the numerals 1, 2, 3, 4 and 5 may be carbon or nitrogen, with the proviso that the ring is aromatic and contains at least 3 carbon atoms;

n is 5 and R^1 at each occurrence is independently selected from halogen, nitro, R^2 , $NR^2_{(m)}$, and OR^2 wherein $m = 0, 1, 2$, or 3 and R^2 at each occurrence is independently selected from H, C_1 - C_8 alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C_3 - C_7 cycloalkyl, C_6 - C_9 polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxyl, alkylcarboxylate, carboxylate, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C_1 - C_4 alkyl and indanyl; wherein the alkyl portion of an R^2 group may be optionally substituted with one or more fluorine atoms, hydroxyl or

4

C₁-C₈alkoxy; and the aryl portion of an R² group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen.

In another aspect, the benzylated PDE4 inhibitor comprises at least one benzyl group (Bzl) of the formula




5

wherein each of the numerals 1, 2, 3, 4 and 5 may be carbon or nitrogen, with the proviso that the ring is aromatic and contains at least 4 carbon atoms;

n is 5 and R¹ at each occurrence is independently selected from halogen, nitro, R², and OR² wherein R² at each occurrence is independently selected from H, C₁-C₈alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C₃-C₇cycloalkyl, C₆-C₉polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, phenyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R² group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-C₈alkoxy; and the phenyl portion of an R² group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen.

In another aspect, the benzylated PDE4 inhibitor comprises at least one benzyl group (Bzl) of the formula




wherein each of the numerals 1, 2, 3, 4 and 5 may be carbon or nitrogen, with the proviso that the ring is aromatic and contains at least 3 carbon atoms;

n is 5 and R¹ at each occurrence is independently selected from halogen, nitro, R², NR²_(m), and OR² wherein m = 0,1,2, or 3 and R² at each occurrence is independently selected from H, C₁-C₈alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C₃-C₇cycloalkyl, C₆-C₉polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl


portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R² group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-C₈alkoxy; and the aryl portion of an R² group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen; with the proviso that either (a) at least one R¹ is OR², or (b) the benzyl ring contains at least two R¹ groups that are not hydrogen.

In another aspect, the benzylated PDE4 inhibitor comprises at least one benzyl group (Bzl) of the formula



wherein each of the numerals 1, 2, 3, 4 and 5 is carbon; n is 5 and R¹ at each occurrence is independently selected from halogen, nitro, R², NR²_(m), and OR² wherein m = 0,1,2, or 3 and R² at each occurrence is independently selected from H, C₁-C₈alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C₃-C₇cycloalkyl, C₆-C₉polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R² group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-C₈alkoxy; and the aryl portion of an R² group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen; with the proviso that R₁ is hydrogen at either 1, 2, or 3 occurrences of n, but not 4 or 5 occurrences.


In one aspect of the present invention, the benzylated PDE4 inhibitor comprises a benzyl group, and independently of the benzyl group the benzylated PDE4 inhibitor further comprises a phenyl group of the formula



wherein p is 5; and R³ at each occurrence is independently selected from halogen, nitro, R⁴, NR⁴_(q), and OR⁴ wherein q = 0,1,2, or 3 and R⁴ at each occurrence is independently selected from H, C₁-C₈alkyl, alkoxyalkyl having 3-7 carbons in the

alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C₃-C₇cycloalkyl, C₆-C₉polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R⁴ group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-C₈alkoxy; and the aryl portion of an R⁴ group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen. In one aspect of the invention R³ is OR⁴ in at least one occurrence wherein R⁴ is selected from C₁-C₈alkyl, C₃-C₇cycloalkyl, and C₆-C₉polycycloalkyl.


In one aspect of the present invention, the phenyl group has the formula



wherein R⁷ is selected from methyl, ethyl, difluoromethyl and trifluoromethyl; and R⁸ is selected from C₃-C₈alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C₃-C₇cycloalkyl, C₆-C₉polycycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R² group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-C₄alkoxy; and the aryl portion (e.g., phenyl) of an R² group may be optionally substituted with C₁-C₄alkyl, C₁-C₄alkoxy or halogen. Optionally, R⁸ is not cyclopentyl.


In one aspect, the PDE4 inhibitor is ROLIPRAMTM or an analog thereof.

For instance, the benzylated ROLIPRAMTM or analog thereof may have the formula:




and isolated stereoisomers, salts and solvates thereof, wherein Bzl is a benzyl group; A is selected from a direct bond, optionally substituted C₁-C₅alkylene, optionally substituted C₂-C₅alkenyl and optionally substituted phenylene; p is 5; and R³ at each occurrence is independently selected from halogen, nitro, R⁴, NR⁴_(q), and OR⁴ wherein q = 0,1,2, or 3 and R⁴ at each occurrence is independently selected from H, C₁-C₈alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C₃-C₇cycloalkyl, C₆-C₉polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R⁴ group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-C₈alkoxy; and the aryl portion of an R⁴ group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen. Optionally, A is a direct bond; and in at least one occurrence, R³ is OR⁴ wherein R⁴ is selected from C₁-C₈alkyl, C₃-C₇cycloalkyl, and C₆-C₉polycycloalkyl.


For instance, the benzylated ROLIPRAM™ or analog thereof may have the formula:




In another aspect the PDE4 inhibitor is ARIFLO™ or an analog thereof. For instance, the benzylated ARIFLO™ or analog thereof may have the formula




- and isolated stereoisomers, salts and solvates thereof, wherein Bzl is a
- 5 benzyl group; p is 5; and R^3 at each occurrence is independently selected from halogen, nitro, R^4 , $NR^4_{(q)}$, and OR^4 wherein $q = 0, 1, 2$, or 3 and R^4 at each occurrence is independently selected from H, C_1 - C_8 alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C_3 - C_7 cycloalkyl, C_6 - C_9 polycycloalkyl, alkylcycloalkyl, hydroxyalkyl,
- 10 carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C_1 - C_4 alkyl and indanyl; wherein the alkyl portion of an R^4 group may be optionally substituted with one or more fluorine atoms, hydroxyl or
- 15 C_1 - C_8 alkoxy; and the aryl portion of an R^4 group may be optionally substituted with C_1 - C_8 alkyl, C_1 - C_8 alkoxy or halogen; R^5 is selected from H, halogen, cyano, C_1 - C_8 alkyl, and C_1 - C_8 alkoxy; and R^6 is selected from H, positively charged species, and C_1 - C_8 alkyl. Optionally, in at least one occurrence, R^3 is OR^4 wherein R^4 is selected from C_1 - C_8 alkyl, C_3 - C_7 cycloalkyl, and C_5 - C_9 bicycloalkyl; and R^6 is selected from H and
- 20 positively charged species. For instance, the benzylated ARIFLO™ or analog thereof may have the formula



As another example, the benzylated ARIFLO™ or analog thereof may have the formula




- and isolated stereoisomers, salts and solvates thereof, wherein Bzl is a benzyl group; p is 5; and R^3 at each occurrence is independently selected from halogen, nitro, R^4 , $NR^4_{(q)}$, and OR^4 wherein $q = 0, 1, 2$, or 3 and R^4 at each occurrence is independently selected from H, C_1 - C_8 alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C_3 - C_7 cycloalkyl, C_6 - C_9 polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C_1 - C_4 alkyl and indanyl; wherein the alkyl portion of an R^4 group may be optionally substituted with one or more fluorine atoms, hydroxyl or C_1 - C_8 alkoxy; and the aryl portion of an R^4 group may be optionally substituted with C_1 - C_8 alkyl, C_1 - C_8 alkoxy or halogen; R^5 is selected from H, halogen, cyano, C_1 - C_8 alkyl, and C_1 - C_8 alkoxy; and R^6 is selected from H, positively charged species, and C_1 - C_8 alkyl. Optionally, in at least one occurrence, R^3 is OR^4 wherein R^4 is selected from C_1 - C_8 alkyl, C_3 - C_7 cycloalkyl, and C_5 - C_9 bicycloalkyl; and R^6 is selected from H and positively charged species. For instance, the benzylated ARIFLO™ or analog thereof may have the formula



In another aspect, the present invention provides a method for improving the therapeutic ratio of PDE4 inhibitors through reduction or elimination of emetogenic

properties while allowing retention of PDE activity, where the method comprises benzylation of a PDE4 inhibitor, wherein the benzylation places a benzyl group (Bzl) on a nitrogen atom of a PDE4 inhibitor, to provide a benzylated PDE4 inhibitor, wherein the benzylated PDE4 inhibitor comprises at least one benzyl group (Bzl) of the
 5 formula




wherein each of the numerals 1, 2, 3, 4 and 5 may be carbon or nitrogen, with the proviso that the ring is aromatic and contains at least 3 carbon atoms; n is 5 and R¹ at each occurrence is independently selected from halogen, nitro, R², NR²_(m), and
 10 OR² wherein m = 0,1,2, or 3 and R² at each occurrence is independently selected from H, C₁-C₈alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C₃-C₇cycloalkyl, C₆-C₉polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl,
 15 heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R² group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-C₈alkoxy; and the aryl portion of an R² group may be optionally substituted with C₁-
 20 C₈alkyl, C₁-C₈alkoxy or halogen; with the proviso that either (a) at least one R¹ is OR², or (b) the benzyl ring contains at least two R¹ groups that are not hydrogen. This may be termed the N-benylation approach to modifying a PDE4 inhibitor. In one aspect, each of the numerals 1, 2, 3, 4 and 5 is carbon. In another aspect, n is 5 and R¹ at each occurrence is independently selected from halogen, nitro, R², NR²_(m), and OR² wherein
 25 m = 0,1,2, or 3 and R² at each occurrence is independently selected from H, C₁-C₈alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C₃-C₇cycloalkyl, C₆-C₉polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, phenyl, phenylalkyl having 1-8 carbons in
 30 the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R² group may be optionally substituted with one or more fluorine atoms,

hydroxyl or C₁-C₈alkoxy; and the phenyl portion of an R² group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen.

The N-benylation method may be applied to a PDE4 inhibitor containing a nitrogen atom, preferably an amide group where the nitrogen atom of the amide group undergoes a benzylation reaction to place a benzyl group on that nitrogen atom. Suitable PDE4 inhibitors for the N-benylation reaction of the present invention include, without limitation, ROLIPRAM™, WAY-PDA-641, Ro-20-1724, RP 73401, CP-80,633-A, AROFYLLINE™ and CIPAMFYLLINE™. The benzylation may replace an extractable hydrogen of the PDE4 inhibitor with a benzyl group. For instance, the PDE4 inhibitor may be reacted with a base in order to extract the extractable hydrogen, followed by reaction with a benzylating agent. In one aspect, the PDE4 inhibitor has an amide group and the extractable hydrogen is located on the nitrogen atom of the amide group.


In one aspect, the benzylated ROLIPRAM™ or analog thereof has the formula:




and isolated stereoisomers, salts and solvates thereof, wherein Bzl is a benzyl group; A is selected from a direct bond, optionally substituted C₁-C₅alkylene, optionally substituted C₂-C₅alkenyl and optionally substituted phenylene; p is 5 and R³ at each occurrence is independently selected from halogen, nitro, R⁴, NR⁴_(q), and OR⁴ wherein q = 0,1,2, or 3 and R⁴ at each occurrence is independently selected from H, C₁-C₈alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C₃-C₇cycloalkyl, C₆-C₉polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R⁴ group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-

12

C₈alkoxy; and the aryl portion of an R⁴ group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen. Optionally, A is a direct bond; p is 1, 2, or 3; and in at least one occurrence, R³ is OR⁴ wherein R⁴ is selected from C₁-C₈alkyl, C₃-C₇cycloalkyl, and C₆-C₉polycycloalkyl. For instance, the benzylated ROLIPRAM™ or analog thereof may have the formula



In another aspect of the inventive method for N-benylation, the PDE4 inhibitor is WAY-PDA-641 or an analog thereof. For instance, the benzylated WAY-PDA-641 or analog thereof may have the formula:




10


and isolated stereoisomers, salts and solvates thereof, wherein Bzl is a benzyl group; p is 5 and R³ at each occurrence is independently selected from halogen, nitro, R⁴, NR⁴_(q), and OR⁴ wherein q = 0,1,2, or 3 and R⁴ at each occurrence is independently selected from H, C₁-C₈alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C₃-C₇cycloalkyl, C₆-C₉polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R⁴ group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-C₈alkoxy; and the aryl portion of an R⁴ group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen. Optionally, p is 1, 2, or 3; and in at least one

13

occurrence, R^3 is OR^4 wherein R^4 is selected from C_1 - C_8 alkyl, C_3 - C_7 cycloalkyl, and C_6 - C_9 polycycloalkyl. For instance, the benzylated WAY-PDA-641 or analog thereof may have the formula:




- 5 In another aspect of the inventive method for N-benylation of a PDE4 inhibitor, the PDE4 inhibitor is Ro-20-1724 or an analog thereof. In one aspect, the benzylated Ro-20-1724 or analog thereof may have the formula:




- and isolated stereoisomers, salts and solvates thereof, wherein Bzl is a
 10 benzyl group; p is 5 and R^3 at each occurrence is independently selected from halogen, nitro, R^4 , $NR^4_{(q)}$, and OR^4 wherein $q = 0, 1, 2$, or 3 and R^4 at each occurrence is independently selected from H, C_1 - C_8 alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C_3 - C_7 cycloalkyl, C_6 - C_9 polycycloalkyl, alkylcycloalkyl, hydroxyalkyl,
 15 carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C_1 - C_4 alkyl and indanyl; wherein the alkyl portion of an R^4 group may be optionally substituted with one or more fluorine atoms, hydroxyl or
 20 C_1 - C_8 alkoxy; and the aryl portion of an R^4 group may be optionally substituted with C_1 - C_8 alkyl, C_1 - C_8 alkoxy or halogen. Optionally, p is 1, 2, or 3; and in at least one occurrence, R^3 is OR^4 wherein R^4 is selected from C_1 - C_8 alkyl, C_3 - C_7 cycloalkyl, and

- 4 -

C₆-C₉polycycloalkyl. For instance, the N-benzylated Ro-20-1724 or analog thereof may have the formula:




In another aspect, the N-benzylated Ro-20-1724 or analog thereof has
5 the formula:




and isolated stereoisomers, salts and solvates thereof, wherein Bzl is a benzyl group; p is 5 and R³ at each occurrence is independently selected from halogen, nitro, R⁴, NR⁴_(q), and OR⁴ wherein q = 0,1,2, or 3 and R⁴ at each occurrence is
10 independently selected from H, C₁-C₈alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C₃-C₇cycloalkyl, C₆-C₉polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl
15 portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R⁴ group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-C₈alkoxy; and the aryl portion of an R⁴ group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen. Optionally, p is 1, 2, or 3; and in at least one
20 occurrence, R³ is OR⁴ wherein R⁴ is selected from C₁-C₈alkyl, C₃-C₇cycloalkyl, and

C₆-C₉polycycloalkyl. For instance, the benzylated Ro-20-1724 or analog thereof may have the formula:




In another aspect, the N-benylation reaction of the present invention employs RP 73401 or an analog thereof as the PDE4 inhibitor. For example, the benzylated RP 73401 or analog thereof may have the formula:




and isolated stereoisomers, salts and solvates thereof, wherein Bzl is a benzyl group; p is 5 and R³ at each occurrence is independently selected from halogen, nitro, R⁴, NR⁴_(q), and OR⁴ wherein q = 0,1,2, or 3 and R⁴ at each occurrence is independently selected from H, C₁-C₈alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C₃-C₇cycloalkyl, C₆-C₉polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R⁴ group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-C₈alkoxy; and the aryl portion of an R⁴ group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen. Optionally, p is 1, 2, or 3; and in at least one

occurrence, R^3 is OR^4 wherein R^4 is selected from C_1 - C_8 alkyl, C_3 - C_7 cycloalkyl, and C_6 - C_9 polycycloalkyl. For instance, the benzylated RP 73401 or analog thereof may have the formula:




- 5 In another aspect of the inventive method for N-benylation of a PDE4 inhibitor, the PDE4 inhibitor is CP-80,633-A or an analog thereof. For example, the benzylated CP-80,633-A or analog thereof may have the formula:




- and isolated stereoisomers, salts and solvates thereof, wherein Bzl is a benzyl group; p is 5 and R^3 at each occurrence is independently selected from halogen, nitro, R^4 , NR^4_q , and OR^4 wherein $q = 0, 1, 2$, or 3 and R^4 at each occurrence is independently selected from H, C_1 - C_8 alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C_3 - C_7 cycloalkyl, C_6 - C_9 polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C_1 - C_4 alkyl and indanyl; wherein the alkyl portion of an R^4 group may be optionally substituted with one or more fluorine atoms, hydroxyl or

C₁-C₈alkoxy; and the aryl portion of an R⁴ group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen. Optionally, p is 1, 2, or 3; and in at least one occurrence, R³ is OR⁴ wherein R⁴ is selected from C₁-C₈alkyl, C₃-C₇cycloalkyl, and C₆-C₉polycycloalkyl. For instance, the benzylated CP-80,633-A or analog thereof may have the formula:



In another aspect of the inventive method for N-benylation of a PDE4 inhibitor, the PDE4 inhibitor is AROFYLLINE™ or an analog thereof. For example, the benzylated AROFYLLINE™ or analog thereof may have the formulae:



10

or


and isolated stereoisomers, salts and solvates thereof, wherein Bzl is a benzyl group; p is 5 and R³ at each occurrence is independently selected from halogen, nitro, R⁴, NR⁴_(q), and OR⁴ wherein q = 0,1,2, or 3 and R⁴ at each occurrence is independently selected from H, C₁-C₈alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C₃-C₇cycloalkyl, C₆-C₉polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an

15


20

18

R⁴ group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-C₈alkoxy; and the aryl portion of an R⁴ group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen. Optionally, p is 1, 2 or 3; and in at least one occurrence, R³ is halogen or OR⁴ wherein R⁴ is selected from C₁-C₈alkyl, C₃-C₇cycloalkyl, and C₆-C₉polycycloalkyl. For example, the benzylated AROFYLLINE™ or analog thereof may have the formula




As another example the benzylated AROFYLLINE™ or analog thereof may have the formula




10


More specific examples of benzylated AROFYLLINE™ or analog thereof compounds prepared by the present invention have the formula:



As another specific example the benzylated AROFYLLINE™ or analog thereof may have the formula:



- 5 In another aspect of the inventive method for N-benylation of a PDE4 inhibitor, the PDE4 inhibitor is CIPAMFYLLINE™ or an analog thereof. For example, the benzylated CIPAMFYLLINE™ or analog thereof may have the formulae:




and isolated stereoisomers, salts and solvates thereof, wherein Bzl is a
10 benzyl group.

In addition the above methods for improving the therapeutic ratio of PDE4 inhibitors through reduction or elimination of emetogenic properties while


allowing retention of PDE activity, the present invention provides PDE4 inhibitors having benzyl groups.

In one aspect, the present invention provides a C-benzylated ROLIPRAM™ or analog thereof compound of the formula




5

and isolated stereoisomers, salts and solvates thereof, wherein Bzl is a benzyl group of the formula




- wherein each of the numerals 1, 2, 3, 4 and 5 may be carbon or nitrogen,
- with the proviso that the ring is aromatic and contains at least 3 carbon atoms; n is 5 and R¹ at each occurrence is independently selected from halogen, nitro, R², NR²_(m), and OR² wherein m = 0,1,2, or 3 and R² at each occurrence is independently selected from H, C₁-C₈alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C₃-C₇cycloalkyl, C₆-C₉polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R² group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-C₈alkoxy; and the aryl portion of an R² group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen; A is selected from a direct bond, optionally substituted C₁-C₅alkylene, optionally substituted C₂-C₅alkenyl and optionally substituted phenylene; p is 5 and R³ at each occurrence is independently selected from halogen, nitro, R⁴, NR⁴_(q), and OR⁴ wherein q = 0,1,2, or 3 and R⁴ at each occurrence is independently selected from H, C₁-C₈alkyl, alkoxyalkyl having 3-7 carbons in the


alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C₃-C₇cycloalkyl, C₆-C₉polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl
 5 portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R⁴ group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-C₈alkoxy; and the aryl portion of an R⁴ group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen. Optionally, A is a direct bond; and in at least one
 10 occurrence, R³ is OR⁴ wherein R² is selected from C₁-C₈alkyl, C₃-C₇cycloalkyl, and C₆-C₉polycycloalkyl. For example, the present invention provides a compound having the formula




In another aspect the present invention provides an N-benzylated
 15 ROLIPRAMTM or analog thereof compound of the formula




and isolated stereoisomers, salts and solvates thereof, wherein Bzl is a benzyl group of the formula




- wherein each of the numerals 1, 2, 3, 4 and 5 may be carbon or nitrogen, with the proviso that the ring is aromatic and contains at least 3 carbon atoms; n is 5 and R¹ at each occurrence is independently selected from halogen, nitro, R², NR²_(m), and OR² wherein m = 0,1,2, or 3 and R² at each occurrence is independently selected from H, C₁-C₈alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C₃-C₇cycloalkyl, C₆-C₉polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R² group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-C₈alkoxy; and the aryl portion of an R² group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen; with the proviso that at least one R¹ is OR²; A is selected from a direct bond, optionally substituted C₁-C₅alkylene, optionally substituted C₂-C₅alkenyl and optionally substituted phenylene; p is 5 and R³ at each occurrence is independently selected from halogen, nitro, R⁴, NR⁴_(q), and OR⁴ wherein q = 0,1,2, or 3 and R⁴ at each occurrence is independently selected from H, C₁-C₈alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C₃-C₇cycloalkyl, C₆-C₉polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R⁴ group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-C₈alkoxy; and the aryl portion of an R⁴ group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen. Optionally, A is a direct bond; and in at least one occurrence, R³ is OR⁴ wherein R⁴ is selected from C₁-C₈alkyl, C₃-C₇cycloalkyl, and C₆-C₉polycycloalkyl. For instance, the present invention provides a compound having the formula



In another aspect the present invention provides a benzylated ARIFLO™ or analog thereof compound of the formulae



5 and isolated stereoisomers, salts and solvates thereof, wherein Bzl is a benzyl group of the formula



wherein each of the numerals 1, 2, 3, 4 and 5 may be carbon or nitrogen, with the proviso that the ring is aromatic and contains at least 3 carbon atoms; n is 5 and


10 R^1 at each occurrence is independently selected from halogen, nitro, R^2 , $NR^2_{(m)}$, and OR^2 wherein $m = 0, 1, 2$, or 3 and R^2 at each occurrence is independently selected from H, C_1 - C_8 alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C_3 - C_7 cycloalkyl, C_6 - C_9 polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate,

15 alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C_1 - C_4 alkyl and indanyl; wherein the alkyl portion of an R^2 group may be optionally substituted with one or more fluorine atoms, hydroxyl or C_1 -

20 C_8 alkoxy; and the aryl portion of an R^2 group may be optionally substituted with C_1 -


C₈alkyl, C₁-C₈alkoxy or halogen; p is 5; and R³ at each occurrence is independently selected from halogen, nitro, R⁴, NR⁴_(q), and OR⁴ wherein q = 0,1,2, or 3 and R⁴ at each occurrence is independently selected from H, C₁-C₈alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C₃-C₇cycloalkyl, C₆-C₉polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R⁴ group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-C₈alkoxy; and the aryl portion of an R⁴ group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen; R⁵ is selected from H, halogen, cyano, C₁-C₈alkyl, and C₁-C₈alkoxy; and R⁶ is selected from H, positively charged species, and C₁-C₈alkyl. Optionally, in at least one occurrence, R³ is OR⁴ wherein R⁴ is selected from C₁-C₈alkyl, C₃-C₇cycloalkyl, and C₅-C₉bicycloalkyl; and R⁶ is selected from H and positively charged species.

For instance, the present invention provides compounds having the formulae




In another aspect the present invention provides a benzylated WAY-PDA-641 or analog thereof compound of the formula:

25




and isolated stereoisomers, salts and solvates thereof, wherein Bzl is a benzyl group of the formula



- 5 wherein each of the numerals 1, 2, 3, 4 and 5 may be carbon or nitrogen, with the proviso that the ring is aromatic and contains at least 3 carbon atoms; n is 5 and R¹ at each occurrence is independently selected from halogen, nitro, R², NR²_(m), and OR² wherein m = 0,1,2, or 3 and R² at each occurrence is independently selected from H, C₁-C₈alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C₃-C₇cycloalkyl, C₆-C₉polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R² group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-C₈alkoxy; and the aryl portion of an R² group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen; p is 5; and R³ at each occurrence is independently selected from halogen, nitro, R⁴, NR⁴_(q), and OR⁴ wherein q = 0,1,2, or 3 and R⁴ at each occurrence is independently selected from H, C₁-C₈alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C₃-C₇cycloalkyl, C₆-C₉polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R⁴ group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-C₈alkoxy; and the aryl portion of an R⁴ group may be optionally


46

substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen. Optionally, in at least one occurrence, R³ is OR⁴ wherein R⁴ is selected from C₁-C₈alkyl, C₃-C₇cycloalkyl, and C₆-C₉polycycloalkyl. For instance, the present invention provides compound having the formula:




5

In another aspect the present invention provides a benzylated Ro-20-1724 or analog thereof compound of the formulae:



or


and isolated stereoisomers, salts and solvates thereof, wherein Bzl is a benzyl group of the formula




wherein each of the numerals 1, 2, 3, 4 and 5 may be carbon or nitrogen, with the proviso that the ring is aromatic and contains at least 3 carbon atoms; n is 5 and R¹ at each occurrence is independently selected from halogen, nitro, R², NR²_(m), and OR² wherein m = 0,1,2, or 3 and R² at each occurrence is independently selected from H, C₁-C₈alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C₃-C₇cycloalkyl, C₆-C₉polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl,

15


- heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R² group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-C₈alkoxy; and the aryl portion of an R² group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen; p is 5; and R³ at each occurrence is independently selected from halogen, nitro, R⁴, NR⁴_(q), and OR⁴ wherein q = 0,1,2, or 3 and R⁴ at each occurrence is independently selected from H, C₁-C₈alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C₃-C₇cycloalkyl, C₆-C₉polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R⁴ group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-C₈alkoxy; and the aryl portion of an R⁴ group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen. Optionally, in at least one occurrence, R³ is OR⁴ wherein R⁴ is selected from C₁-C₈alkyl, C₃-C₇cycloalkyl, and C₅-C₉bicycloalkyl. For example, the present invention provides a compound having the formulae




28



In another aspect the present invention provides a benzylated RP 73401 or analog thereof compound of the formula:




- 5 and isolated stereoisomers, salts and solvates thereof, wherein Bzl is a benzyl group of the formula




- wherein each of the numerals 1, 2, 3, 4 and 5 may be carbon or nitrogen, with the proviso that the ring is aromatic and contains at least 3 carbon atoms; n is 5 and
- 10 R^1 at each occurrence is independently selected from halogen, nitro, R^2 , $NR^2_{(m)}$, and OR^2 wherein $m = 0, 1, 2$, or 3 and R^2 at each occurrence is independently selected from H, C_1 - C_8 alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C_3 - C_7 cycloalkyl, C_6 - C_9 polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate,
- 15 alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be

optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R² group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-C₈alkoxy; and the aryl portion of an R² group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen; p is 5; and R³ at each occurrence is independently selected from halogen, nitro, R⁴, NR⁴_(q), and OR⁴ wherein q = 0,1,2, or 3 and R⁴ at each occurrence is independently selected from H, C₁-C₈alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C₃-C₇cycloalkyl, C₆-C₉polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R⁴ group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-C₈alkoxy; and the aryl portion of an R⁴ group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen. Optionally, in at least one occurrence, R³ is OR⁴ wherein R⁴ is selected from C₁-C₈alkyl, C₃-C₇cycloalkyl, and C₆-C₉polycycloalkyl. For example, the present invention provides a compound having the formula:




20

In another aspect the present invention provides a benzylated CP-80,633-A or analog thereof compound of the formula:




and isolated stereoisomers, salts and solvates thereof, wherein Bzl is a benzyl group of the formula




- 5 wherein each of the numerals 1, 2, 3, 4 and 5 may be carbon or nitrogen, with the proviso that the ring is aromatic and contains at least 3 carbon atoms; n is 5 and R^1 at each occurrence is independently selected from halogen, nitro, R^2 , $NR^2_{(m)}$, and OR^2 wherein $m = 0, 1, 2$, or 3 and R^2 at each occurrence is independently selected from H, C_1 - C_8 alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C_3 - C_7 cycloalkyl, C_6 - C_9 polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be
- 10 optionally substituted with C_1 - C_4 alkyl and indanyl; wherein the alkyl portion of an R^2 group may be optionally substituted with one or more fluorine atoms, hydroxyl or C_1 - C_8 alkoxy; and the aryl portion of an R^2 group may be optionally substituted with C_1 - C_8 alkyl, C_1 - C_8 alkoxy or halogen; p is 5; and R^3 at each occurrence is independently selected from halogen, nitro, R^4 , $NR^4_{(q)}$, and OR^4 wherein $q = 0, 1, 2$, or 3 and R^4 at each
- 15 occurrence is independently selected from H, C_1 - C_8 alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C_3 - C_7 cycloalkyl, C_6 - C_9 polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8
- 20 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C_1 - C_4 alkyl and indanyl; wherein the alkyl portion of an R^4 group may be optionally substituted with one or more fluorine
- 25


atoms, hydroxyl or C₁-C₈alkoxy; and the aryl portion of an R⁴ group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen. Optionally, in at least one occurrence, R³ is OR⁴ wherein R⁴ is selected from C₁-C₈alkyl, C₃-C₇cycloalkyl, and C₆-C₉polycycloalkyl. For example, the present invention provides a compound having
 5 the formula:



In another aspect the present invention provides a benzylated AROFYLLINE™ or analog thereof of the formulae:




10 and isolated stereoisomers, salts and solvates thereof, wherein Bzl is a benzyl group of the formula




wherein each of the numerals 1, 2, 3, 4 and 5 may be carbon or nitrogen, with the proviso that the ring is aromatic and contains at least 3 carbon atoms; n is 5 and
 15 R¹ at each occurrence is independently selected from halogen, nitro, R², NR²_(m), and OR² wherein m = 0,1,2, or 3 and R² at each occurrence is independently selected from H, C₁-C₈alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in


- the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C₃-C₇cycloalkyl, C₆-C₉polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion,
- 5 phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R² group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-C₈alkoxy; and the aryl portion of an R² group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen; p is 5; and R³ at each occurrence is independently
- 10 selected from halogen, nitro, R⁴, NR⁴_(q), and OR⁴ wherein q = 0,1,2, or 3 and R⁴ at each occurrence is independently selected from H, C₁-C₈alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C₃-C₇cycloalkyl, C₆-C₉polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide,
- 15 alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R⁴ group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-C₈alkoxy; and the aryl portion of an R⁴ group may be optionally
- 20 substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen. Optionally, in at least one occurrence, R³ is halogen or OR⁴ wherein R⁴ is selected from C₁-C₈alkyl, C₃-C₇cycloalkyl, and C₆-C₉polycycloalkyl. For example, the present invention provides a benzylated AROFYLLINE™ or analog thereof compound having the formulae:




33



In another aspect the present invention provides a benzylated CIPAMFYLLINE™ or analog thereof compound of the formulae:



5 and isolated stereoisomers, salts and solvates thereof, wherein Bzl is a benzyl group of the formula




wherein each of the numerals 1, 2, 3, 4 and 5 may be carbon or nitrogen, with the proviso that the ring is aromatic and contains at least 3 carbon atoms; n is 5 and

10 R^1 at each occurrence is independently selected from halogen, nitro, R^2 , $NR^2_{(m)}$, and OR^2 wherein $m = 0, 1, 2$, or 3 and R^2 at each occurrence is independently selected from H, C_1 - C_8 alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C_3 - C_7 cycloalkyl, C_6 - C_9 polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl,


15 heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C_1 - C_4 alkyl and indanyl; wherein the alkyl portion of an R^2

group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-C₈alkoxy; and the aryl portion of an R² group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen. For example, the present invention provides a benzylated CIPAMFYLLINE™ or analog thereof compound of the formula:



5

As another example, the present invention provides a benzylated CIPAMFYLLINE™ or analog thereof compound of the formula:



In addition to the above-mentioned method for improving the therapeutic ratio of PDE4 inhibitors through reduction or elimination of emetogenic properties while allowing retention of PDE activity, and the compounds of the present invention, the present invention also provides for inventive pharmaceutical compositions comprising a compound of the present invention and a pharmaceutically acceptable carrier, diluent or excipient. Furthermore, the present invention provides various methods for treating and/or preventing various biological, e.g., medical, conditions, where those methods employ a compound of the present invention, or a pharmaceutical composition of the present invention. Those methods for treating and/or preventing various biological conditions include the following.

A method for treating or preventing an inflammatory condition or disease in a patient, comprising administering to the patient in need thereof an amount of a benzylated PDE4 inhibitor, where the amount is effective to treat or prevent the inflammatory condition or disease of the patient, and where the benzylated PDE4 inhibitor is prepared by a benzylation method described herein. In addition, the present

invention provides a method for treating or preventing an inflammatory condition or disease in a patient, comprising administering to the patient in need thereof an amount of a benzylated compound as described herein, where the amount is effective to treat or prevent the inflammatory condition or disease of the patient. In one aspect, the PDE4 inhibitor inhibits an enzyme selected from phosphodiesterase 4 A, B, C or D or any combination thereof including all splice variants of PDE4 A, B, C and D. In various aspects, the inflammatory condition or disease is an autoimmune condition or disease; or involves acute or chronic inflammation of bone and/or cartilage compartments of joints; or is an arthritis selected from rheumatoid arthritis, gouty arthritis or juvenile rheumatoid arthritis; or is asthma; or is associated with the dysregulation of T-cells; or the inflammatory condition or disease is multiple sclerosis; or the inflammatory condition or disease is pulmonary sarcoidosis; or the inflammatory condition or disease is ocular inflammation or allergy; or the inflammatory condition or disease is an inflammatory bowel disease, where the inflammatory bowel disease may be Crohn's disease or ulcerative colitis; or the inflammatory condition or disease is an inflammatory cutaneous disease, where the inflammatory cutaneous disease may be psoriasis or dermatitis; or the inflammatory condition or disease is chronic obstructive pulmonary disease (COPD), bronchitis, emphysema or acute respiratory distress syndrome (ARDS). In another aspect, the inflammatory condition or disease is associated with elevated levels of inflammatory cytokines. In various aspects, the inflammatory cytokine is IL-2, IL-4 or IL-5; or the inflammatory cytokine is IFN- γ ; or the inflammatory cytokine is TNF- α .

A method for treating or preventing a disease or condition in a patient, where the disease or condition is associated with pathological conditions that are modulated by inhibiting enzymes associated with secondary cellular messengers, the method comprising administering to the patient in need thereof an amount of a benzylated PDE4 inhibitor, wherein the amount is effective to treat or prevent a disease or condition associated with pathological conditions that are modulated by inhibiting enzymes associated with secondary cellular messengers, and wherein the benzylated PDE4 inhibitor is prepared by a benzylation process according to the present invention. In addition, the present invention provides a method for treating or preventing a disease or condition in a patient, where the disease or condition is associated with pathological conditions that are modulated by inhibiting enzymes associated with secondary cellular messengers, the method comprising administering to the patient in need thereof an amount of a benzylated compound according to the present invention, or a pharmaceutical composition containing a benzylated compound according to the

present invention, wherein the amount is effective to treat or prevent a disease or condition associated with pathological conditions that are modulated by inhibiting enzymes associated with secondary cellular messengers. In various aspects, the enzyme is a cyclic AMP phosphodiesterase; or the enzyme is phosphodiesterase 4.

5 A method of treating or preventing transplant rejection in a patient, the method comprising administering to the patient in need thereof an amount of a benzylated PDE4 inhibitor, where the amount is effective to treat or prevent transplant rejection in the patient, and wherein the benzylated PDE4 inhibitor is prepared by a benzylation process according to the present invention. In addition, the present invention
10 provides a method of treating or preventing transplant rejection in a patient, the method comprising administering to the patient in need thereof an amount of a benzylated compound according to the present invention, or a composition comprising a compound according to the present invention and a pharmaceutically acceptable carrier, diluent and excipient, where the amount is effective to treat or prevent transplant rejection in
15 the patient. In one aspect the rejection is due to graft versus host disease.

 A method of treating or preventing uncontrolled cellular proliferation in a patient, the method comprising administering to the patient in need thereof an amount of a benzylated PDE4 inhibitor, where the amount is effective to treat or prevent uncontrolled cellular proliferation in the patient, and wherein the benzylated PDE4
20 inhibitor is prepared by a process according to the present invention. In addition, the present invention provides a method of treating or preventing uncontrolled cellular proliferation in a patient, the method comprising administering to the patient in need thereof an amount of a benzylated compound according to the present invention, or a composition comprising a benzylated compound according to the present invention and
25 a pharmaceutically acceptable carrier, diluent or excipient, where the amount is effective to treat or prevent uncontrolled cellular proliferation in the patient. In one aspect, the uncontrolled cellular proliferation is caused by a cancer selected from leukemia and solid tumors.

 A method of treating or preventing conditions associated with the central
30 nervous system (CNS) in a patient, the method comprising administering to the patient in need thereof an amount of a benzylated PDE4 inhibitor, where the amount is effective to treat or prevent conditions associated with the central nervous system (CNS) in the patient, and wherein the benzylated PDE4 inhibitor is prepared by a process according to the present invention. In addition, the present invention provides a
35 method of treating or preventing conditions associated with the central nervous system (CNS) in a patient, the method comprising administering to the patient in need thereof

an amount of a benzylated compound according to the present invention, or a composition comprising a benzylated compound according to the present invention and a pharmaceutically acceptable carrier, diluent or excipient, where the amount is effective to treat or prevent conditions associated with the central nervous system (CNS) in the patient. In one aspect, the condition associated with the central nervous system (CNS) is depression. In another aspect, the condition is long-term memory ostentation and learning enhancement.

A method of treating or preventing diseases in a patient, the disease associated with viral infection, the method comprising administering to the patient in need thereof an amount of a benzylated PDE4 inhibitor, where the amount is effective to treat or prevent diseases associated with viral infection in the patient, and wherein the benzylated PDE4 inhibitor is prepared by a process according to the present invention. In addition, the present invention provides a method of treating or preventing diseases in a patient, the disease associated with viral infection, the method comprising administering to the patient in need thereof an amount of a benzylated compound according to the present invention, or a composition comprising a pharmaceutically acceptable carrier, diluent or excipient in combination with a benzylated compound of the present invention, where the amount is effective to treat or prevent diseases associated with viral infection in the patient. In one aspect, the viral infection is due to the human immunodeficiency virus (HIV) and the disease is acquired immunodeficiency syndrome (AIDS).

A method of treating or preventing diseases in a patient, the disease associated with infection by a parasite, the method comprising administering to the patient in need thereof an amount of a benzylated PDE4 inhibitor, where the amount is effective to treat or prevent diseases associated with infection of the patient by a parasite, and wherein the benzylated PDE4 inhibitor is prepared by a process according to the present invention. In another aspect, the present invention provides a method of treating or preventing diseases in a patient, the disease associated with infection by a parasite, the method comprising administering to the patient in need thereof an amount of a benzylated compound according to the present invention, or a composition comprising a pharmaceutically acceptable carrier, diluent or excipient in combination with a benzylated compound of the present invention, where the amount is effective to treat or prevent diseases associated with infection of the patient by a parasite. In one aspect, the parasitic infection is due to the trypanosome Bruce and the disease is African sleeping sickness disease.

20


A method of treating or preventing cystic fibrosis in a patient, the method comprising administering to the patient in need thereof an amount of a benzylated PDE4 inhibitor, where the amount is effective to treat or prevent cystic fibrosis in the patient, where the amount is effective to treat or prevent cystic fibrosis, and wherein the benzylated PDE4 inhibitor is prepared by a benzylation process according to the present invention. In addition, the present invention provides a method of treating or preventing cystic fibrosis in a patient, the method comprising administering to the patient in need thereof an amount of a benzylated compound according to the present invention, or a composition comprising a pharmaceutically acceptable carrier, diluent or excipient in combination with a benzylated compound of the present invention, where the amount is effective to treat or prevent cystic fibrosis.

DETAILED DESCRIPTION OF THE INVENTION

It has been surprisingly discovered that compounds exhibiting PDE4 inhibitory activity and emetogenic effects will exhibit reduced emetogenic effects when the compound contains certain chemical substitution, and in particular certain chemical substitution including a benzyl group.

Definitions


As used herein, and unless otherwise specified, a benzyl group (Bzl, sometimes referred to as Bn) refers to a group of the formula



20

wherein each of the numerals 1, 2, 3, 4 and 5 may be carbon or nitrogen, and when a numeral is carbon then that numeral may be substituted with an organic or inorganic group.

In one aspect, a benzyl group (Bzl) refers to a group of the formula




25

wherein each of the numerals 1, 2, 3, 4 and 5 may be carbon or nitrogen, with the proviso that the ring is aromatic and contains at least 3 carbon atoms;

n is 5 and R^1 at each occurrence is independently selected from halogen, nitro, R^2 , $NR^2_{(m)}$, and OR^2 wherein $m = 0, 1, 2$, or 3 and R^2 at each occurrence is independently selected from H, C_1 - C_8 alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C_3 - C_7 cycloalkyl, C_6 - C_9 polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C_1 - C_4 alkyl and indanyl; wherein the alkyl portion of an R^2 group may be optionally substituted with one or more fluorine atoms, hydroxyl or C_1 - C_8 alkoxy; and the aryl portion of an R^2 group may be optionally substituted with C_1 - C_8 alkyl, C_1 - C_8 alkoxy or halogen.


In one aspect, a benzyl group (Bzl) refers to a group of the formula



wherein each of the numerals 1, 2, 3, 4 and 5 may be carbon or nitrogen, with the proviso that the ring is aromatic and contains at least 4 carbon atoms;

n is 5 and R^1 at each occurrence is independently selected from halogen, nitro, R^2 , and OR^2 wherein and R^2 at each occurrence is independently selected from H, C_1 - C_8 alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C_3 - C_7 cycloalkyl, C_6 - C_9 polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, phenyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C_1 - C_4 alkyl and indanyl; wherein the alkyl portion of an R^2 group may be optionally substituted with one or more fluorine atoms, hydroxyl or C_1 - C_8 alkoxy; and the phenyl portion of an R^2 group may be optionally substituted with C_1 - C_8 alkyl, C_1 - C_8 alkoxy or halogen.


In one aspect, a benzyl group (Bzl) refers to a group of the formula



wherein each of the numerals 1, 2, 3, 4 and 5 may be carbon or nitrogen, with the proviso that the ring is aromatic and contains at least 3 carbon atoms;

n is 5 and R^1 at each occurrence is independently selected from halogen, nitro, R^2 , $NR^2_{(m)}$, and OR^2 wherein $m = 0, 1, 2$, or 3 and R^2 at each occurrence is independently selected from H, C_1 - C_8 alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C_3 - C_7 cycloalkyl, C_6 - C_9 polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C_1 - C_4 alkyl and indanyl; wherein the alkyl portion of an R^2 group may be optionally substituted with one or more fluorine atoms, hydroxyl or C_1 - C_8 alkoxy; and the aryl portion of an R^2 group may be optionally substituted with C_1 - C_8 alkyl, C_1 - C_8 alkoxy or halogen; with the proviso that at least one R^1 is OR^2 .


In one aspect, a benzyl group (Bzl) refers to a group of the formula



wherein each of the numerals 1, 2, 3, 4 and 5 is carbon;

n is 5 and R^1 at each occurrence is independently selected from halogen, nitro, R^2 , $NR^2_{(m)}$, and OR^2 wherein $m = 0, 1, 2$, or 3 and R^2 at each occurrence is independently selected from H, C_1 - C_8 alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C_3 - C_7 cycloalkyl, C_6 - C_9 polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C_1 - C_4 alkyl and indanyl; wherein the alkyl portion of an R^2 group may be optionally substituted with one or more fluorine atoms, hydroxyl or C_1 - C_8 alkoxy; and the aryl portion of an R^2 group may be optionally substituted with C_1 - C_8 alkyl, C_1 - C_8 alkoxy or halogen; with the proviso that at least one R^1 is halogen.

In one embodiment, each of the aromatic ring atoms of the benzyl group is a carbon atom, so that the benzyl group has the formula



wherein n is 0, 1, 2, 3 or 4, and independently at each occurrence,

R¹ is selected from halogen, nitro, R², NR²_(m), and OR² wherein m = 0, 1, 2, or 3 and R² is independently selected from H, C₁-C₈alkyl, alkoxyalkyl having 3-7
 5 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C₃-C₇cycloalkyl, C₆-C₉polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8
 10 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R² group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-C₈alkoxy; and the aryl portion of an R² group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen.

In any of the above embodiments, in a further embodiment of the present
 15 invention, n is 1, 2 or 3 and in at least one occurrence R¹ is halogen.

In a compound as disclosed herein, when any variable occurs more than one time in any constituent or in a benzylated compound, its definition on each occurrence is independent of its definition at every other occurrence. Combinations of substituents and/or variables are permissible only if such combinations result in stable
 20 compounds. The compounds useful in the methods and compositions of the present invention, as well as the compounds of the present invention, may have asymmetric centers and occur as racemates, racemic mixtures and as individual diastereomers, or enantiomers with all isomeric forms being included in the present invention. A racemate or racemic mixture does not imply a 50:50 mixture of stereoisomers. Also,
 25 unless otherwise indicated, the terms "a" and "an" refer to one, or more than one, of the indicated items. For example, "a compound" includes one and more than one compound, and "an" isomer refers to one and more than one isomer.

In the benzylated compounds, a salt is preferably a pharmaceutically acceptable salt, where salts includes acid addition salts and base addition salts. A
 30 "pharmaceutically acceptable salt" and "salts thereof" means organic or inorganic salts of the pharmaceutically important molecule. A pharmaceutically acceptable salt may involve the inclusion of another molecule such as an acetate ion, a succinate ion or other counterion. The counterion may be any organic or inorganic moiety that stabilizes the charge on the parent compound. Furthermore, a pharmaceutically

important organic molecule may have more than one charged atom in its structure. Situations where multiple charged atoms are part of the molecule may have multiple counterions. Hence, the molecule of a pharmaceutically acceptable salt may contain one or more than one charged atoms and may also contain, one or more than one
5 counterion. The desired charge distribution is determined according to methods of drug administration. Examples of pharmaceutically acceptable salts are well known in the art but, without limiting the scope of the present invention, exemplary presentations can be found in the Physician's Desk Reference, The Merck Index, The Pharmacopoeia and Goodman & Gilman's The Pharmacological Basis of Therapeutics.

10 Acid addition salts refer to those salts formed from benzylated compounds of the present invention and inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid and the like, and/or organic acids such as acetic acid, propionic acid, glycolic acid, pyruvic acid, oxalic acid, maleic acid, malonic acid, succinic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, cinnamic acid, mandelic
15 acid, methanesulfonic acid, ethanesulfonic acid, p-toluenesulfonic acid, salicylic acid and the like.

Base addition salts include those salts derived from benzylated compounds of the present invention and inorganic bases such as sodium, potassium, lithium, ammonium, calcium, magnesium, iron, zinc, copper, manganese, aluminum
20 salts and the like. Suitable salts include the ammonium, potassium, sodium, calcium and magnesium salts derived from pharmaceutically acceptable organic non-toxic bases include salts of primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines and basic ion exchange resins, such as isopropylamine, trimethylamine, diethylamine, triethylamine, tripropylamine,
25 ethanolamine, 2-dimethylaminoethanol, 2-diethylaminoethanol, trimethylamine, dicyclohexylamine, lysine, arginine, histidine, caffeine, procaines, hydrabamine, choline, betaine, ethylenediamine, glucosamine, methylglucamine, theobromine, purines, piperazine, piperidine, N-ethylpiperidine, and the like.

30 "Alkoxy" refers to alkyl-O-, where "O" represents oxygen and alkyl is defined below.

"Alkoxyalkyl" refers to alkyl¹-O-alkyl²-, where "O" represents oxygen and alkyl¹ represents the alkoxy portion, *i.e.*, alkyl¹-O is alkoxy, while alkyl² represents the alkyl portion of the alkoxyalkyl group. In an alkoxyalkyl group, each of alkyl¹ and alkyl² is an alkyl group as defined below, with the comment that alkyl¹ is monovalent as it has
35 one open valence site that is bound to oxygen, while alkyl² is divalent, as it is bound to both oxygen and the remainder of the molecule.

"Alkyl" refers to an acyclic chain of carbon atoms that may be branched or unbranched (linear). Methyl, ethyl, propyl (including *n*-propyl and *iso*-propyl) butyl (including *n*-butyl, *iso*-butyl, *sec*-butyl, and *t*-butyl), pentyl (including numerous isomers) and hexyl (including numerous isomers) are alkyl groups having 1 to 6 carbon atoms (commonly referred to as lower alkyl groups), and are exemplary of alkyl groups of the invention. As referred to herein, an alkyl group may have unsaturation between any two carbons, *i.e.*, the alkyl group may have double and/or triple bonds. Thus, the term alkyl encompasses alkenyl groups, which are unsaturated aliphatic groups having at least one double bond, as well as alkynyl groups, which are unsaturated aliphatic groups which may be either straight- or branched-chain and have one or more triple bonds. In various embodiments the alkyl groups have no more than about 20 carbons atoms, or no more than about 18, or 16, or 14, or 12, or 10 carbons atoms, and may be ethyl, propynyl, 4-methylpentynyl and so on, and structural isomers thereof. In addition, the term alkyl encompasses monovalent as well as polyvalent alkyl groups, where methyl (CH₃-) is an exemplary monovalent alkyl group, while methylene (-CH₂-) is an exemplary polyvalent, and more specifically a divalent alkyl group. In one embodiment, alkyl excludes methyl. In another embodiment, the alkyl group has at least 3 carbon atoms.


"Alkylamide" refers to an amide group wherein either or both of the carbon and nitrogen atoms are bonded to alkyl group(s).

"Alkylcarboxyl" refers to a divalent alkyl group wherein one of the open valence sites is bonded to a carboxyl group, while the remaining open valence site is bonded to the remainder of the molecule.

"Alkylcarboxylate" refers to a divalent alkyl group wherein one of the open valence sites is bonded to a carboxylate group, while the remaining open valence site is bonded to the remainder of the molecule.

"Alkylcycloalkyl" refers to a cycloalkyl group that is bonded both to an alkyl group and to the remainder of the molecule. Each of alkyl and cycloalkyl are defined herein.


"Amide" refers to a group of the formula



where the amide group may be joined to the remainder of the molecule through either the carbonyl group or the nitrogen atom.

"Aryl" refers to an aromatic hydrocarbon ring system. The ring system may be monocyclic or fused polycyclic (*e.g.*, bicyclic, tricyclic, etc.). In various embodiments, the monocyclic aryl ring is C5-C10, or C5-C7, or C5-C6, where these carbon numbers refer to the number of carbon atoms that make up the ring system. A C6 ring system, *i.e.*, a phenyl ring, is a preferred aryl ring. In various embodiments, the polycyclic ring is a bicyclic aryl ring, where preferred bicyclic aryl rings are C8-C12, or C9-C10. A naphthyl ring, which has 10 carbon atoms, is a preferred polycyclic aryl ring. Aryl rings may be substituted or unsubstituted. In one embodiment, the aryl ring is unsubstituted. In another embodiment, the aryl ring is substituted with 1 substituent (*i.e.*, the aryl ring is monosubstituted), or 1-2 substituents, or 1-3 substituents, or 1-4 substituents, etc.

"Carboxylate" refers to a group of the formula



i.e., -COO⁻, where there is a positively charged counterion present.

- 15 "Carboxyl" or "Carboxylic acid" refers to a protonated carboxylate.
- "Cycloalkyl" refers to a cyclic alkyl group. Suitable cycloalkyl groups include, for example, cyclohexyl, cyclopropyl, cyclopentyl, and cycloheptyl. Cycloalkenyl is an example of a cycloalkyl group, where suitable cycloalkenyl groups include, for example, cyclopentenyl and cyclohexenyl.
- 20 "Halogen" refers to fluorine, chlorine, bromine and iodine.
- "Heteroalkyl" refers to a saturated or unsaturated, straight or branched, chain containing carbon and at least one heteroatom. The heteroalkyl group may, in various embodiments, have one heteroatom, or 1-2 heteroatoms, or 1-3 heteroatoms, or 1-4 heteroatoms. Heteroalkyl chains may contain from 1 to 18 (*i.e.*, 1-18) member atoms (carbon and heteroatoms) in the chain, and in various embodiments contain 1-12, or 1-6, or 1-4 member atoms. Independently, in various embodiments, the heteroalkyl group has zero branches (*i.e.*, is a straight chain), one branch, two branches, or more than two branches. Independently, in one embodiment, the heteroalkyl group is saturated. In another embodiment, the heteroalkyl group is unsaturated. In various
- 25 atoms (carbon and heteroatoms) in the chain, and in various embodiments contain 1-12, or 1-6, or 1-4 member atoms. Independently, in various embodiments, the heteroalkyl group has zero branches (*i.e.*, is a straight chain), one branch, two branches, or more than two branches. Independently, in one embodiment, the heteroalkyl group is saturated. In another embodiment, the heteroalkyl group is unsaturated. In various
- 30 embodiments, the unsaturated heteroalkyl may have one double bond, two double bonds, more than two double bonds, and/or one triple bond, two triple bonds, or more than two triple bonds. Heteroalkyl chains may be substituted or unsubstituted. In one embodiment, the heteroalkyl chain is unsubstituted. In another embodiment, the heteroalkyl chain is substituted. A substituted heteroalkyl chain may have 1 substituent

(i.e., be monosubstituted), or may have 1-2 substituents, or 1-3 substituents, or 1-4 substituents, etc. Exemplary heteroalkyl groups include esters (-C(=O)-OR) and ketones (-C(=O)-). In another embodiment, the heteroalkyl group is not substituted.

"Heteroaryl" is an aromatic ring system or a semi-aromatic system of
5 rings or a pseudo aromatic ring or rings containing carbon and at least one heteroatom in at least one of the rings. The heteroaryl group may, in various embodiments, have one heteroatom, or 1-2 heteroatoms, or 1-3 heteroatoms, or 1-4 heteroatoms in the ring. The heteroaryl group may further include more than one ring system, which in various
10 embodiments may include one heteroatom or 1-2 heteroatoms, or 1-3 heteroatoms, or 1 heteroatom in each ring system, or 1-4 heteroatoms in each ring system. The heteroaryl group which comprises more than one ring system may, in various embodiments have one or more than one of the ring systems aromatic. Heteroaryl rings may be monocyclic or polycyclic, where the polycyclic ring may contain fused, spiro or bridged
15 ring junctions. In one embodiment, the heteroaryl is selected from monocyclic and bicyclic. Monocyclic heteroaryl rings may contain from about 5 to about 10 member atoms (carbon and heteroatoms), preferably from 5-7, and most preferably from 5-6 member atoms in the ring. Bicyclic heteroaryl rings may contain from about 8-12 member atoms, or 9-10 member atoms in the ring. The heteroaryl ring may be unsubstituted or substituted. In one embodiment, the heteroaryl ring is unsubstituted.
20 In another embodiment, the heteroaryl ring is substituted. The substituted heteroaryl ring may contain 1 substituent, or 1-2 substituents, or 1-3 substituents, or 1-4 substituents, etc. Exemplary heteroaryl rings include benzofuran, benzothiophene, furan, imidazole, indole, isothiazole, oxazole, pyrazine, pyrazole, pyridazine, pyridine, pyrimidine, pyrrole, quinoline, thiazole and thiophene. In one embodiment, the heteroaryl group does not
25 have any substituents.

"Heteroatom" is a nitrogen, sulfur, oxygen or silicon atom. Groups containing more than one heteroatom may contain different heteroatoms.


"Heterocycloalkyl" refers to a saturated or unsaturated, monocyclic or polycyclic (e.g., bicyclic, tricyclic, etc.) ring containing carbon and at least one
30 heteroatom. Heterocyclic aliphatic rings are not aromatic *per se* but may be pseudo-aromatic and/or readily be made aromatic through methods known in the art. The heterocyclic aliphatic ring may, in various embodiments, have one heteroatom, or 1-2 heteroatoms, or 1-3 heteroatoms, or 1-4 heteroatoms, etc. In one embodiment, the heterocyclic aliphatic ring is monocyclic, where the monocyclic ring may have 3-10, or
35 4-7, or 5-6 member atoms. In another embodiment, the heterocyclic aliphatic ring is polycyclic, where in various embodiments, the ring may be bicyclic, or may be

tricyclic, or may be either bicyclic or tricyclic. A polycyclic ring system may have one or more fused, spiro or bridged ring systems. The polycyclic heterocyclic aliphatic ring system may have 6-12, or 9-10 member atoms. The heterocyclic ring may be unsubstituted or substituted. In one embodiment, the heterocyclic ring is unsubstituted.

5 In another embodiment, the heterocyclic ring is substituted. The substituted heterocyclic ring may contain 1 substituent, or 1-2 substituents, or 1-3 substituents, or 1-4 substituents, etc. Exemplary heterocyclic aliphatic rings include piperazyl, morpholiny, tetrahydrofuranyl, tetrahydropyranyl and piperidyl. In another embodiment, the heterocycloalkyl is not substituted.

10 "Hydroxyalkyl" refers to an alkyl group that is bonded to both a hydroxy (HO-) group and to the remainder of the molecule. In other words, a group that may be represented as HO-alkyl-.

"Indanyl" refers to an indane group, *i.e.*, a group of the structure



15 wherein one of the carbons of the indane group is bonded to the remainder of the molecule.

"Phenoxyalkyl" refers to phenyl-O-alkyl-, where "phenyl-O" is the phenoxy portion of the phenoxyalkyl group, and "alkyl" is the alkyl portion of the phenoxyalkyl group. The alkyl portion of the phenoxyalkyl group is defined above,
20 with the comment that in a phenoxyalkyl radical, the alkyl group is divalent.

"Phenylalkyl" refers to phenyl-alkyl-, where "phenyl" is the phenyl portion of the phenylalkyl group, and "alkyl" is the alkyl portion of the phenylalkyl group. The alkyl portion of the phenylalkyl group is defined above, with the comment that in a phenylalkyl radical, the alkyl group is divalent.

25 "Polycycloalkyl" refers to an arrangement of carbon atoms wherein at least one carbon atom is a part of at least two separately identifiable rings. The polycycloalkyl group may contain bridging between two carbon atoms, where bicyclo[1.1.0]butyl, bicyclo[3.2.1]octyl, bicyclo[5.2.0]nonyl, tricycl[2.2.1.0¹]heptyl, norbornyl and pinanyl are representative examples. The polycycloalkyl group may
30 contain one or more fused ring systems, where decaliny (radical from decalin) and perhydroanthracenyl are representative examples. The polycycloalkyl group may contain a spiro union, in which a single atom is the only common member of two rings. Spiro[3.4]octyl, spiro[3.3]heptyl and spiro[4.5]decyl are representative examples.

Polycycloalkyl groups having 6, 7, 8 or 9 carbon atoms, *i.e.*, C₆-C₉polycycloalkyl, are a preferred polycycloalkyl group of the present invention.

In describing the present invention, the following abbreviations may be used, where these abbreviations have the meaning as indicated in the following Table.

5

Abbreviation	Full name
5-ASA	5-aminosalicylic acid
Ab	Antibody
ABTS	2,2'-azino-di-[3-ethylbenzthiazoline sulphonate]
ACD	Acid citrate dextrose
AcOH	Acetic Acid
ACVP	American College of Veterinary Practice
ANOVA	Analysis of Variance
Ar	Argon
BCR-ABL	Oncogene in chromosome 9:22 translocation in CML
BINAP	2,2'-Bis(diphenylphosphino)-1,1'-binaphthyl
Bn	Benzyl
BnBr	Benzyl Bromide
BOC	tert-Butoxycarbonyl
cAMP	Cyclic adenosine 3' -5'-monophosphate
cat	Catalytic
CD	Cluster designation
CFA	Complete Freund's adjuvant
cGMP	Cyclic guanosine 3'-5'-monophosphate
CIA	Collagen Induced Arthritis
CLL	Chronic lymphocytic leukemia
CML	Chronic myelogenous leukemia
CNS	Central Nervous System
Con A	Concanavalin A
COX	Cyclooxygenase
cPent	Cyclopentyl
cPentBr	Cyclopentyl bromide
CRE	cAMP response element
CsA	Cyclosporin A
DMAP	4-Dimethylaminopyridine
DMARD	Disease modifying anti-rheumatic drug

Abbreviation	Full name
DMF	<i>N,N</i> -Dimethylformamide
DMSO	dimethylsulfoxide
DNA	Deoxyriboneucleic acid
DNBS	2,4-Dinitrobenzene sulphoric acid
dppf	1,1'-Bis(diphenylphosphino)ferrocene
dppp	1,3-Bis(diphenylphosphino)propane
EC ₅₀	Concentration at which a 50% of maximum observable effect is noted
EDTA	Ethylenediaminetetraacetic acid
ELISA	Enzyme-linked immunosorbent assay
EtOAc	Ethyl acetate
EtOH	Ethyl alcohol
FBS	Fetal bovine serum
FCS	Fetal calf serum
fMLP	Formyl-methionyl leucine phenylalanine
g.i.	Gastrointestinal
H & E	Haematoxylin and eosin
HARBS	High affinity rolipram binding site
HBSS	Hanks Balanced Salt Solution
HMPA	Hexamethylphosphoramide
HPLC	High pressure liquid chromatography
i.p.	intraperitoneal
IBD	Inflammatory bowel disease
IBMX	3-isobutyl-1-methylxanthine
IC	Inhibitory concentration
IC ₅₀	Concentration at which 50% inhibition is observed
IFA	Incomplete Freund's adjuvant
IFN- γ	Interferon gamma
IL	Interleukin
LAH	Lithium aluminum hydride
LDA	Lithium diisopropylamide
LN	Lymph node
LPS	lipopolysaccharide
LTB ₄	Leukotriene B ₄
luc	luciferase


Abbreviation	Full name
Me	Methyl
MeOH	Methyl alcohol
MHC	Major histocompatibility class
MLR	Mixed lymphocyte reaction
MPO	myeloperoxidase
Ms	Methanesulfonyl
MsCl	Methanesulfonyl chloride
NBS	<i>N</i> -Bromosuccinimide
<i>n</i> -BuLi	<i>n</i> -Butyllithium
<i>n</i> -BuSH	<i>n</i> -Butanethiol
NF- κ B	Nuclear factor kappa B
NSAID	Non-steroidal anti-inflammatory drug
p.t.	Post-transplant
PBS	Phosphate buffered saline
Pcc	Pigeon cytochrome C
PDE	Phosphodiesterase
PEG	Polyethylene glycol
PG	Prostaglandin
PMS	Phenazine methosulfate
PMSF	Phenyl methyl sulfonyl fluoride
pTsOH	<i>p</i> -Toluenesulfonic acid monohydrate
Py	Pyridine
RA	Rheumatoid arthritis
RF	Rheumatoid factor
R _f	Retardation factor
ROS	Reactive oxygen species
RPMI	Rosewell Park Memorial Institute
RTX	Resiniferitoxin
SAR	Structure activity relationship
TBAF	Tetrabutylammonium fluoride
TBDMS	<i>tert</i> -Butyldimethylsilyl
TBDMSCl	<i>tert</i> -Butyldimethylsilyl chloride
TCR	T-cell receptor
TEA	Triethylamine
Tf	Trifluoromethanesulfonyl

Abbreviation	Full name
TFA	Trifluoroacetic acid
Th	T helper
THF	Tetrahydrofuran
TNBS	Trinitrobenzene sulfonic acid
TNF- α	Tumour necrosis factor alpha
Trolox [®]	6-hydroxy-2,5,7,8-tetramethylchroman-2-carboxylic acid
TsOH	<i>p</i> -Toluenesulfonic acid monohydrate
XTT	2,3-bis[2-methoxy-4-nitro-5-sulfo-phenyl]-2H-tetrazolium 5-carboxanilide inner salt
μ M	Micro molar

Method of the Present Invention

In one aspect, the present invention is directed to a method for improving the therapeutic ratio of PDE4 inhibitors through reduction or elimination of emetogenic properties while allowing retention of PDE activity. The method comprises benzylation of a PDE4 inhibitor, wherein the benzylation places a benzyl group on a carbon (referred to herein as C-benylation) or nitrogen (referred to herein as N-benylation) atom of a PDE4 inhibitor, to provide a benzylated PDE4 inhibitor. In one aspect of the invention, the benzylation places a benzyl group on a carbon atom of a PDE4 inhibitor. In another aspect of the invention, the benzylation places a benzyl group on a nitrogen atom of a PDE4 inhibitor. In one aspect of the invention, the benzylation places a benzyl group on a carbon atom or a nitrogen atom of a PDE4 inhibitor. In another aspect, the benzylation places two benzyl groups on a PDE4 inhibitor.

The addition of a benzyl group to a PDE4 inhibitor is readily accomplished by reacting the PDE4 inhibitor (or a protected version thereof) with a benzylating agent (or a protected version thereof) under conditions such that the benzylating agent forms a covalent bond to a carbon or nitrogen atom of the PDE4 inhibitor. As used herein, a benzylating agent includes a benzyl group as defined herein having a leaving group attached to the methylene carbon of the benzyl group, *i.e.*, a leaving group alpha to the aromatic ring. Suitable leaving groups include halogens, *e.g.*, chloride and bromide, and sulfonates, *e.g.*, mesylate and tosylate. Such a benzylating agent has the general formula as shown below.



wherein n is 0, 1, 2, 3 or 4, and independently at each occurrence,

R^1 is selected from halogen, nitro, R^2 , $NR^2_{(m)}$, and OR^2 wherein $m = 0, 1, 2$, or 3 and R^2 is independently selected from H, C_1 - C_8 alkyl, alkoxyalkyl having 3-7
 5 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C_3 - C_7 cycloalkyl, C_6 - C_9 polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8
 10 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C_1 - C_4 alkyl and indanyl; wherein the alkyl portion of an R^2 group may be optionally substituted with one or more fluorine atoms, hydroxyl or C_1 - C_8 alkoxy; and the aryl portion of an R^2 group may be optionally substituted with C_1 - C_8 alkyl, C_1 - C_8 alkoxy or halogen.


Reaction conditions whereby a PDE4 inhibitor will react with a
 15 benzylating agent so as to form a benzylated PDE4 inhibitor include basic conditions. In the presence of basic conditions, the one or more extractable hydrogens of the PDE4 inhibitor will react with base so that the PDE4 inhibitor forms the corresponding nucleophilic anion. The nucleophilic center will then react with the benzylating agent with concomitant displacement of the leaving group, so as to join the PDE4 inhibitor to
 20 the benzyl group. Benzylation chemistry in general is well known in the art, and may be applied to the goal of benzylating PDE4 inhibitors according to the present invention.

The basic conditions are created by combining the PDE4 inhibitor with a base, typically in a suitable solvent. Suitable bases must be selected based on the
 25 acidity of the hydrogen atom being removed from the PDE4 inhibitor. That is, a relatively more basic compound must be employed when the hydrogen being removed from the PDE4 inhibitor is relatively less extractable. Suitable bases for most instances include sodium hydride, potassium hydride, metal amine salts such as lithium amine salts, *e.g.*, lithium diisopropylamide, and sodium amine salts, lithium hydroxide,
 30 sodium hydroxide, potassium hydroxide, sodium alkoxide or potassium alkoxide where alkoxide refers to O-alkyl and alkyl is defined herein.

Scheme 1 depicts general synthetic methodology that may be used for the introduction of a benzyl group at a location alpha to a carbonyl group. Scheme 1 depicts this methodology with an ester-containing molecule, and more specifically with

a lactone, however, the same methodology may be used when the carbonyl group is part of a lactam or other amide-containing molecule, or when the carbonyl group is part of a non-lactone ester group or ketone group or carboxylic acid group. As discussed in more detail below, this general methodology may be applied to a carbonyl-containing compound that has substitution and/or when the benzylating agent has substitution.


Scheme 1



Although Scheme 1 illustrates one methodology that may be employed according to the present invention using an unsubstituted benzylation agent (and, more specifically, benzyl bromide, abbreviated as BnBr) to afford a benzylated product, any number of substituted benzylation agents can be used to generate a benzylated product with various substitution patterns on the benzyl ring. Substituted benzyl bromides are available commercially or may be generated from the corresponding substituted benzyl alcohol, benzaldehyde, benzoic acid or benzoic ester.

For example, substituted benzyl bromides can be prepared as outlined in Scheme 2. Thus, treatment of commercially available benzyl alcohol 1 with benzyl bromide and potassium carbonate in toluene gives the corresponding benzyloxy derivative 2, which is treated, without purification, with PBr₃ in diethyl ether to give the desired bromide compound 3 in quantitative yield. Any number of compounds related to compound 3 could be produced using similar methodology but starting with a different substituted benzyl alcohol.

Scheme 2




Substituted benzyl bromide compounds can also be prepared from commercially available substituted benzaldehydes, benzoic acids and benzoic esters by first converting these compounds to the corresponding alcohol. Benzyl aldehydes may be reduced to the corresponding alcohol by standard reducing conditions, *e.g.*, sodium borohydride in methanol. In some instances, it may be necessary to protect other functionality on the benzylaldehyde compound in order that the other functionality is not undesirably modified by the reduction conditions.

10 It may be necessary to protect certain reactive groups present in the PDE4 inhibitor from the benzylation reaction. Suitable protecting groups and methods for their use are described in "Protective Groups in Organic Synthesis", 3rd Edition, by Greene, T.W. and Wuts, P.G.M., John Wiley & Sons, 1999.

As for selection of the PDE4 inhibitor, many of these are known in the art and may be advantageously benzylated according to the present invention. Suitable PDE4 inhibitors include, without limitation, ROLIPRAM™, ARIFLO™, WAY-PDA 641, Ro-20-1724, RP 73401, CP-80,633-A, AROFYLLINE™, and CIPAMFYLLINE™. In a preferred embodiment, benzylation of a PDE4 inhibitor occurs by placing a benzyl group on a carbon atom of the PDE4 inhibitor. PDE4 inhibitors suitable for benzylation on a carbon atom include, without limitation, ROLIPRAM™ and ARIFLO™. In another aspect, the present invention employs PDE4 inhibitors suitable for benzylation on a nitrogen atom, where such PDE4 inhibitors include, without limitation, WAY-PDA-641, Ro-20-1724, RP 73401, CP-80,633-A, AROFYLLINE™, and CIPAMFYLLINE™. The following literature references, in addition to those provided in the Background of the present specification, describe various PDE4 inhibitors and methodology to identify a PDE4 inhibitor: PCT International Publication No. WO92/12961; *J. Med. Chem.* 1997, 40, 1417-1421; *Organic Process Research & Development* 1998, 2, 157-168; *J. Med.*

Chem., 1994, 37, 1696-1703; and U.S. Patent Nos. 5,124,455; 5,814,651; 3,636,039; 4,308,278; 5,223,504; 5,734,051.

In one aspect of the present invention, the benzylated PDE4 inhibitor comprises a benzyl group, and independently of the benzyl group the benzylated PDE4 inhibitor further comprises the group




wherein p is 5; and R^3 at each occurrence is independently selected from halogen, nitro, R^4 , $NR^4_{(q)}$, and OR^4 wherein $q = 0, 1, 2$, or 3 and R^4 at each occurrence is independently selected from H, C_1 - C_8 alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C_3 - C_7 cycloalkyl, C_6 - C_9 polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C_1 - C_4 alkyl and indanyl; wherein the alkyl portion of an R^4 group may be optionally substituted with one or more fluorine atoms, hydroxyl or C_1 - C_8 alkoxy; and the aryl portion of an R^4 group may be optionally substituted with C_1 - C_8 alkyl, C_1 - C_8 alkoxy or halogen. In one aspect of the invention R^3 is OR^4 in at least one occurrence wherein R^4 is selected from C_1 - C_8 alkyl, C_3 - C_7 cycloalkyl, and C_6 - C_9 polycycloalkyl.

C-Benzylation of PDE4 Inhibitors

In one aspect of the invention, the benzylation may provide a benzyl group attached to a carbon atom of the PDE4 inhibitor. In various embodiments of this aspect of the invention, the benzyl group being added to the PDE4 inhibitor has one of the following definitions:


A benzyl group (Bzl) of the formula



wherein each of the numerals 1, 2, 3, 4 and 5 may be carbon or nitrogen, with the proviso that the ring is aromatic and contains at least 3 carbon atoms; n is 5 and


R^1 at each occurrence is independently selected from halogen, nitro, R^2 , $NR^2_{(m)}$, and OR^2 wherein $m = 0, 1, 2$, or 3 and R^2 at each occurrence is independently selected from H, C_1 - C_8 alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C_3 - C_7 cycloalkyl, C_6 - C_9 polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxyl, alkylcarboxylate, carboxylate, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1- carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C_1 - C_4 alkyl and indanyl; wherein the alkyl portion of an R^2 group may be optionally substituted with one or more fluorine atoms, hydroxyl or C_1 - C_8 alkoxy; and the aryl portion of an R^2 group may be optionally substituted with C_1 - C_8 alkyl, C_1 - C_8 alkoxy or halogen; or

A benzyl group (Bzl) of the formula




wherein each of the numerals 1, 2, 3, 4 and 5 may be carbon or nitrogen, with the proviso that the ring is aromatic and contains at least 4 carbon atoms; n is 5 and R^1 at each occurrence is independently selected from halogen, nitro, R^2 , and OR^2 wherein and R^2 at each occurrence is independently selected from H, C_1 - C_8 alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C_3 - C_7 cycloalkyl, C_6 - C_9 polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, phenyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C_1 - C_4 alkyl and indanyl; wherein the alkyl portion of an R^2 group may be optionally substituted with one or more fluorine atoms, hydroxyl or C_1 - C_8 alkoxy; and the phenyl portion of an R^2 group may be optionally substituted with C_1 - C_8 alkyl, C_1 - C_8 alkoxy or halogen; or

A benzyl group (Bzl) of the formula



- wherein each of the numerals 1, 2, 3, 4 and 5 may be carbon or nitrogen, with the proviso that the ring is aromatic and contains at least 3 carbon atoms; n is 5 and R^1 at each occurrence is independently selected from halogen, nitro, R^2 , $NR^2_{(m)}$, and OR^2 wherein $m = 0, 1, 2$, or 3 and R^2 at each occurrence is independently selected from
- 5 H, C_1 - C_8 alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C_3 - C_7 cycloalkyl, C_6 - C_9 polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion,
- 10 phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C_1 - C_4 alkyl and indanyl; wherein the alkyl portion of an R^2 group may be optionally substituted with one or more fluorine atoms, hydroxyl or C_1 - C_8 alkoxy; and the aryl portion of an R^2 group may be optionally substituted with C_1 - C_8 alkyl, C_1 - C_8 alkoxy or halogen; with the proviso that either (a) at least one R^1 is OR^2 ,
- 15 or (b) the benzyl ring contains at least two R^1 groups that are not hydrogen.


A benzyl group (Bzl) of the formula




- wherein each of the numerals 1, 2, 3, 4 and 5 may be carbon or nitrogen, with the proviso that the ring is aromatic and contains at least 3 carbon atoms; n is 5 and R^1 at each occurrence is independently selected from halogen, nitro, R^2 , $NR^2_{(m)}$, and OR^2 wherein $m = 0, 1, 2$, or 3 and R^2 at each occurrence is independently selected from
- 20 H, C_1 - C_8 alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C_3 - C_7 cycloalkyl, C_6 - C_9 polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion,
- 25 phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C_1 - C_4 alkyl and indanyl; wherein the alkyl portion of an R^2 group may be optionally substituted with one or more fluorine atoms, hydroxyl or C_1 - C_8 alkoxy; and the aryl portion of an R^2 group may be optionally substituted with C_1 - C_8 alkyl, C_1 - C_8 alkoxy or halogen; with the proviso that R_1 is hydrogen at either 1, 2, or 3 occurrences of n, but not 4 or 5 occurrences.
- 30

-7-


In one aspect of the invention, the PDE4 inhibitor is ROLIPRAM™ or an analog thereof. In one embodiment, the benzylated ROLIPRAM™ or analog thereof according to the inventive method has the formula:



- 5 and isolated stereoisomers, salts and solvates thereof, wherein Bzl is a benzyl group; A is selected from a direct bond, optionally substituted C₁-C₅alkylene, optionally substituted C₂-C₅alkenyl and optionally substituted phenylene; p is 5; and R³ at each occurrence is independently selected from halogen, nitro, R⁴, NR⁴_(q), and OR⁴ wherein q = 0,1,2, or 3 and R⁴ at each occurrence is independently selected from H,
- 10 C₁-C₈alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C₃-C₇cycloalkyl, C₆-C₉polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion,
- 15 phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R⁴ group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-C₈alkoxy; and the aryl portion of an R⁴ group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen. In one aspect of the invention R³ is OR⁴ in at least
- 20 one occurrence wherein R⁴ is selected from C₁-C₈alkyl, C₃-C₇cycloalkyl, and C₆-C₉polycycloalkyl. In another embodiment, A is a direct bond; p is 1, 2, or 3; and in at least one occurrence, R³ is OR⁴ wherein R⁴ is selected from C₁-C₈alkyl, C₃-C₇cycloalkyl, and C₆-C₉polycycloalkyl. In another embodiment, the benzylated ROLIPRAM™ or analog thereof has the formula:




In a different aspect, the PDE4 inhibitor undergoing the benzylation reaction of the present invention is ARIFLO™ or an analog thereof. In one embodiment, the benzylated ARIFLO™ or analog thereof produced the present invention has the formula




and isolated stereoisomers, salts and solvates thereof, wherein Bzl is a benzyl group; p is 5; and R^3 at each occurrence is independently selected from halogen, nitro, R^4 , $NR^4_{(q)}$, and OR^4 wherein $q = 0, 1, 2$, or 3 and R^4 at each occurrence is independently selected from H, C_1 - C_8 alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C_3 - C_7 cycloalkyl, C_6 - C_9 polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C_1 - C_4 alkyl and indanyl; wherein the alkyl portion of an R^4 group may be optionally substituted with one or more fluorine atoms, hydroxyl or C_1 - C_8 alkoxy; and the aryl portion of an R^4 group may be optionally substituted with C_1 - C_8 alkyl, C_1 - C_8 alkoxy or halogen; R^6 is selected from H, positively charged species, and C_1 - C_8 alkyl; and R^5 is selected from H, halogen, nitro, cyano, C_1 - C_8 alkyl, and C_1 - C_8 alkoxy. In one aspect of the invention R^3 is OR^4 in at least one occurrence wherein R^4 is selected from C_1 - C_8 alkyl, C_3 - C_7 cycloalkyl, and C_6 - C_9 polycycloalkyl; R^6 is selected from H, positively charged species, and C_1 - C_8 alkyl; and R^5 is selected from H,

halogen, nitro, cyano, C₁-C₈alkyl, and C₁-C₈alkoxy. In one embodiment, p is 1, 2 or 3; and in at least one occurrence, R³ is OR⁴ wherein R⁴ is selected from C₁-C₈alkyl, C₃-C₇cycloalkyl, and C₅-C₉bicycloalkyl, R⁶ is selected from H and positively charged species; and R⁵ is selected from H, halogen, nitro, cyano, C₁-C₃alkyl, and C₁-C₃alkoxy.

5 In another embodiment, the benzylated ARIFLO™ or analog thereof has the formula




In another embodiment, the benzylated ARIFLO™ or analog thereof has the formula



10 and isolated stereoisomers, salts and solvates thereof, wherein Bzl is a benzyl group; p is 5; and R³ at each occurrence is independently selected from halogen, nitro, R⁴, NR⁴_(q), and OR⁴ wherein q = 0,1,2, or 3 and R⁴ at each occurrence is independently selected from H, C₁-C₈alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C₃-C₇cycloalkyl, C₆-C₉polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, 15 carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R⁴ group may be optionally substituted with one or more fluorine atoms, hydroxyl or 20 C₁-C₈alkoxy; and the aryl portion of an R⁴ group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen. In one embodiment, p is 1, 2 or 3; and in at least one occurrence, R³ is OR⁴ wherein R⁴ is selected from C₁-C₈alkyl, C₃-C₇cycloalkyl, and C₆-C₉polycycloalkyl, R⁶ is selected from H and positively charged species; and R⁵ is

60


selected from H, halogen, cyano, C₁-C₃alkyl, and C₁-C₃alkoxy. In one embodiment, the benzylated ARIFLO™ or analog thereof has the formula



N-Benzylation of PDE4 Inhibitors


- 5 In one aspect of the invention, the benzylation may provide a benzyl group attached to a nitrogen atom of the PDE4 inhibitor. In a preferred embodiment of this aspect of the invention, the benzyl group has the following definition:

A benzyl group (Bzl) of the formula




- 10 wherein each of the numerals 1, 2, 3, 4 and 5 may be carbon or nitrogen, with the proviso that the ring is aromatic and contains at least 3 carbon atoms; n is 5 and R¹ at each occurrence is independently selected from halogen, nitro, R², NR²_(m), and OR² wherein m = 0,1,2, or 3 and R² at each occurrence is independently selected from H, C₁-C₈alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C₃-C₇cycloalkyl, C₆-C₉polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, 15 alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be 20 optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R² group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-C₈alkoxy; and the aryl portion of an R² group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen; with the proviso that either (a) at least one R¹ is OR², or (b) the benzyl ring contains at least two R¹ groups that are not hydrogen.

- 25 A benzyl group (Bzl) of the formula




- wherein each of the numerals 1, 2, 3, 4 and 5 is carbon, n is 5 and R¹ at each occurrence is independently selected from halogen, nitro, R², NR²_(m), and OR² wherein m = 0, 1, 2, or 3 and R² at each occurrence is independently selected from H, C₁-C₈alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C₃-C₇cycloalkyl, C₆-C₉polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R² group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-C₈alkoxy; and the aryl portion of an R² group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen.

- 15 A benzyl group (Bzl) of the formula




- n is 5 and R¹ at each occurrence is independently selected from halogen, nitro, R², NR²_(m), and OR² wherein m = 0, 1, 2, or 3 and R² at each occurrence is independently selected from H, C₁-C₈alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C₃-C₇cycloalkyl, C₆-C₉polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, phenyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R² group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-C₈alkoxy; and the phenyl portion of an R² group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen.

- In another aspect, the benzylated PDE4 inhibitor comprises at least one benzyl group (Bzl) of the formula



- wherein each of the numerals 1, 2, 3, 4 and 5 may be carbon or nitrogen, with the proviso that the ring is aromatic and contains at least 3 carbon atoms; n is 5 and R¹ at each occurrence is independently selected from halogen, nitro, R², NR²_(m), and OR² wherein m = 0,1,2, or 3 and R² at each occurrence is independently selected from H, C₁-C₈alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C₃-C₇cycloalkyl, C₆-C₉polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R² group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-C₈alkoxy; and the aryl portion of an R² group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen; with the proviso that either (a) at least one R¹ is OR², or (b) the benzyl ring contains at least two R¹ groups that are not hydrogen.

In another aspect, the benzylated PDE4 inhibitor comprises at least one benzyl group (Bzl) of the formula




- wherein each of the numerals 1, 2, 3, 4 and 5 is carbon; n is 5 and R¹ at each occurrence is independently selected from halogen, nitro, R², NR²_(m), and OR² wherein m = 0,1,2, or 3 and R² at each occurrence is independently selected from H, C₁-C₈alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C₃-C₇cycloalkyl, C₆-C₉polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R² group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-C₈alkoxy; and the aryl portion of an R² group may be optionally substituted with C₁-

67

C₈alkyl, C₁-C₈alkoxy or halogen; with the proviso that R₁ is hydrogen at either 1, 2, or 3 occurrences of n, but not 4 or 5 occurrences.

In another aspect, the benzylated PDE4 inhibitor comprises at least one benzyl group (Bzl) of the formula




5

wherein each of the numerals 1, 2, 3, 4 and 5 may be carbon or nitrogen, with the proviso that the ring is aromatic and contains at least 3 carbon atoms;

n is 5 and R¹ at each occurrence is independently selected from halogen and OR² wherein R² at each occurrence is independently selected from H and C₁-C₈alkyl; with the provisos that the benzyl ring contain no more than three hydrogen substituents and no more than one methoxy substituent.

In one aspect of the present invention where benzylation provides an N-benzylated PDE4 inhibitor, the PDE4 inhibitor is ROLIPRAM™ or analog thereof and the N-benzylated ROLIPRAM™ has the formula:




15


and isolated stereoisomers, salts and solvates thereof, wherein Bzl is a benzyl group; A is selected from a direct bond, optionally substituted C₁-C₅alkylene, optionally substituted C₂-C₅alkenyl and optionally substituted phenylene; p is 5; and R³ at each occurrence is independently selected from halogen, nitro, R⁴, NR⁴_(q), and OR⁴ wherein q = 0,1,2, or 3 and R⁴ at each occurrence is independently selected from H, C₁-C₈alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C₃-C₇cycloalkyl, C₆-C₉polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be

25

optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R⁴ group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-C₈alkoxy; and the aryl portion of an R⁴ group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen. In one embodiment, A is a direct bond; p is 1, 2, or 3; and in at least one occurrence, R³ is OR⁴ wherein R⁴ is selected from C₁-C₈alkyl, C₃-C₇cycloalkyl, and C₆-C₉polycycloalkyl. In one embodiment, benzylated ROLIPRAM™ or analog thereof has the formula




In one aspect of the present invention where benzylation provides an N-benzylated PDE4 inhibitor, the PDE4 inhibitor is WAY-PDA-641 or an analog thereof. See, e.g., U.S. Patent No. 5,124,455. In one embodiment, the benzylated WAY-PDA-641 or analog thereof has the formula:



and isolated stereoisomers, salts and solvates thereof, wherein Bzl is a benzyl group; p is 5; and R³ at each occurrence is independently selected from halogen, nitro, R⁴, NR⁴_(q), and OR⁴ wherein q = 0,1,2, or 3 and R⁴ at each occurrence is independently selected from H, C₁-C₈alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C₃-C₇cycloalkyl, C₆-C₉polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an


R^4 group may be optionally substituted with one or more fluorine atoms, hydroxyl or C_1 - C_8 alkoxy; and the aryl portion of an R^4 group may be optionally substituted with C_1 - C_8 alkyl, C_1 - C_8 alkoxy or halogen. In one embodiment, p is 1, 2 or 3; and in at least one occurrence, R^3 is OR^4 wherein R^4 is selected from C_1 - C_8 alkyl, C_3 - C_7 cycloalkyl, and

5 C_6 - C_9 polycycloalkyl. In another embodiment, the benzylated WAY-PDA-641 or analog thereof has the formula:



In one aspect of the present invention where benzylation provides an N-benzylated PDE4 inhibitor, the PDE4 inhibitor is Ro-20-1724 or an analog thereof.

10 See, e.g., U.S. Patent Nos. 3,636,039 and 4,308,278. In one embodiment, the benzylated Ro-20-1724 or analog thereof has the formula:




and isolated stereoisomers, salts and solvates thereof, wherein Bzl is a benzyl group; p is 5; and R^3 at each occurrence is independently selected from halogen, nitro, R^4 , $NR^4_{(q)}$, and OR^4 wherein $q = 0, 1, 2$, or 3 and R^4 at each occurrence is

15 independently selected from H, C_1 - C_8 alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C_3 - C_7 cycloalkyl, C_6 - C_9 polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl,


20 heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C_1 - C_4 alkyl and indanyl; wherein the alkyl portion of an

66

R^4 group may be optionally substituted with one or more fluorine atoms, hydroxyl or C_1 - C_8 alkoxy; and the aryl portion of an R^4 group may be optionally substituted with C_1 - C_8 alkyl, C_1 - C_8 alkoxy or halogen. In one embodiment, p is 1, 2 or 3; and in at least one occurrence, R^3 is OR^4 wherein R^4 is selected from C_1 - C_8 alkyl, C_3 - C_7 cycloalkyl, and C_5 - C_9 bicycloalkyl. In one embodiment, the benzylated Ro-20-1724 or analog thereof has the formula:




In another embodiment, the benzylated Ro-20-1724 or analog thereof has the formula:




10

and isolated stereoisomers, salts and solvates thereof, wherein Bzl is a benzyl group; p is 5; and R^3 at each occurrence is independently selected from halogen, nitro, R^4 , $NR^4_{(q)}$, and OR^4 wherein $q = 0, 1, 2$, or 3 and R^4 at each occurrence is independently selected from H, C_1 - C_8 alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C_3 - C_7 cycloalkyl, C_6 - C_9 polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C_1 - C_4 alkyl and indanyl; wherein the alkyl portion of an R^4 group may be optionally substituted with one or more fluorine atoms, hydroxyl or

C₁-C₈alkoxy; and the aryl portion of an R⁴ group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen. In one embodiment, p is 1, 2 or 3; and in at least one occurrence, R³ is OR⁴ wherein R⁴ is selected from C₁-C₈alkyl, C₃-C₇cycloalkyl, and C₆-C₉polycycloalkyl. In another embodiment, the benzylated Ro-20-1724 or analog thereof has the formula:




In one aspect of the present invention where benzylation provides an N-benzylated PDE4 inhibitor, the PDE4 inhibitor is RP 73401 or an analog thereof. See, e.g., PCT International Publication No. 92/12961; Organic Process Research & Development 1998, 2, 157-168; and *J. Med. Chem.* 1994, 37, 1696-1703. In one embodiment, the benzylated RP 73401 or analog thereof has the formula:




and isolated stereoisomers, salts and solvates thereof, wherein Bzl is a benzyl group; p is 5; and R³ at each occurrence is independently selected from halogen, nitro, R⁴, NR⁴_(q), and OR⁴ wherein q = 0,1,2, or 3 and R⁴ at each occurrence is independently selected from H, C₁-C₈alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C₃-C₇cycloalkyl, C₆-C₉polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl

- portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R⁴ group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-C₈alkoxy; and the aryl portion of an R⁴ group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen. In one embodiment, p is 1, 2 or 3; and in at least one occurrence, R³ is OR⁴ wherein R⁴ is selected from C₁-C₈alkyl, C₃-C₇cycloalkyl, and C₆-C₉polycycloalkyl. In another embodiment, the benzylated RP 73401 or analog thereof has the formula:




- 10 In one aspect of the present invention where benzylation provides an N-benzylated PDE4 inhibitor, the PDE4 inhibitor is CP-80,633-A or an analog thereof. *See, e.g.*, U.S. Patent No. 5,814,651. In one embodiment, the benzylated CP-80,633-A or analog thereof has the formula:




- 15 and isolated stereoisomers, salts and solvates thereof, wherein Bzl is a benzyl group; p is 5; and R³ at each occurrence is independently selected from halogen, nitro, R⁴, NR⁴_(q), and OR⁴ wherein q = 0,1,2, or 3 and R⁴ at each occurrence is independently selected from H, C₁-C₈alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in

the alkyl portion, C₃-C₇cycloalkyl, C₆-C₉polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may
 5 be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R⁴ group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-C₈alkoxy; and the aryl portion of an R⁴ group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen. In one embodiment, p is 1, 2 or 3; and in at least one occurrence, R³ is OR⁴ wherein R⁴ is selected from C₁-C₈alkyl, C₃-C₇cycloalkyl, and
 10 C₆-C₉polycycloalkyl. In one embodiment, the benzylated CP-80,633-A or analog thereof has the formula:




A benzylated compound as disclosed herein includes stereoisomers in admixture and isolated form. Shown below are two stereoisomers of benzylated CP-
 15 80,633-A according to the invention.




In one aspect of the present invention where benzylation provides an N-benzylated PDE4 inhibitor, the PDE4 inhibitor is AROFYLLINE™ or an analog

70

thereof. In one embodiment, the benzylated AROFYLLINE™ or analog thereof has the formulae:




- and isolated stereoisomers, salts and solvates thereof, wherein Bzl is a benzyl group; p is 5; and R³ at each occurrence is independently selected from halogen, nitro, R⁴, NR⁴_(q), and OR⁴ wherein q = 0, 1, 2, or 3 and R⁴ at each occurrence is independently selected from H, C₁-C₈alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C₃-C₇cycloalkyl, C₆-C₉polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R⁴ group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-C₈alkoxy; and the aryl portion of an R⁴ group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen. In one embodiment, p is 1, 2 or 3; and in at least one occurrence, R³ is halogen or OR⁴ wherein R⁴ is selected from C₁-C₈alkyl, C₃-C₇cycloalkyl, and C₆-C₉polycycloalkyl. In one embodiment, the benzylated AROFYLLINE™ or analog thereof has the formula:




71

while in another embodiment, the benzylated AROFYLLINE™ or analog thereof has the formula:



In one aspect of the present invention where benzylation provides an N-
 5 benzylated PDE4 inhibitor, the PDE4 inhibitor is CIPAMFYLLINE™ or an analog thereof. In one embodiment, the benzylated CIPAMFYLLINE™ or analog thereof is selected from the formulae:




and

and isolated stereoisomers, salts and solvates thereof, wherein Bzl is a benzyl group.


10 Benzylated PDE4 Inhibitors

In various aspects, the present invention provides the following specific benzylated PDE4 inhibitors.

A C-benzylated ROLIPRAM™ or analog thereof compound of the formula




and isolated stereoisomers, salts and solvates thereof, wherein Bzl is a benzyl group; A is selected from a direct bond, optionally substituted C₁-C₅alkylene, optionally substituted C₂-C₅alkenyl and optionally substituted phenylene; p is 5; and R³ at each occurrence is independently selected from halogen, nitro, R⁴, NR⁴_(q), and OR⁴ wherein q = 0,1,2, or 3 and R⁴ at each occurrence is independently selected from H, C₁-C₈alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C₃-C₇cycloalkyl, C₆-C₉polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R⁴ group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-C₈alkoxy; and the aryl portion of an R⁴ group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen. In a further embodiment, A is a direct bond; p is 1, 2, or 3; and in at least one occurrence, R³ is OR⁴ wherein R⁴ is selected from C₁-C₈alkyl, C₃-C₇cycloalkyl, and C₆-C₉polycycloalkyl. In a preferred embodiment, R³ at the para position relative to A is alkoxy, and R³ at one meta position is selected from alkyl having at least 3 carbons and alkoxy having at least 3 carbons. In a further embodiment, the benzylated PDE4 inhibitor has the formula:




73

A benzylated ARIFLO™ or analog thereof compound of the formulae



and isolated stereoisomers, salts and solvates thereof, wherein Bzl is a benzyl group of the formula




5

- wherein each of the numerals 1, 2, 3, 4 and 5 may be carbon or nitrogen, with the proviso that the ring is aromatic and contains at least 3 carbon atoms; n is 5 and R¹ at each occurrence is independently selected from halogen, nitro, R², NR²_(m), and OR² wherein m = 0,1,2, or 3 and R² at each occurrence is independently selected from
- 10 H, C₁-C₈alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C₃-C₇cycloalkyl, C₆-C₉polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion,
- 15 phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R² group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-C₈alkoxy; and the aryl portion of an R² group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen; p is 5; and R³ at each occurrence is independently
- 20 selected from halogen, nitro, R⁴, NR⁴_(q), and OR⁴ wherein q = 0,1,2, or 3 and R⁴ at each occurrence is independently selected from H, C₁-C₈alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C₃-C₇cycloalkyl, C₆-C₉polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8
- 25 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the


alkyl portion of an R^4 group may be optionally substituted with one or more fluorine atoms, hydroxyl or C_1 - C_8 alkoxy; and the aryl portion of an R^4 group may be optionally substituted with C_1 - C_8 alkyl, C_1 - C_8 alkoxy or halogen; R^5 is selected from H, halogen, cyano, C_1 - C_8 alkyl, and C_1 - C_8 alkoxy; and R^6 is selected from H, positively charged species, and C_1 - C_8 alkyl. Optionally, in at least one occurrence, R^3 is OR^4 wherein R^4 is selected from C_1 - C_8 alkyl, C_3 - C_7 cycloalkyl, and C_5 - C_9 bicycloalkyl; and R^6 is selected from H and positively charged species.

In one aspect, the benzylated ARIFLO™ or analog thereof compound has a formula,




10

In another aspect, the benzylated ARIFLO™ or analog thereof compound has a formula,




In another aspect, the benzylated ARIFLO™ or analog thereof compound has a formula,

15




75


In another aspect, the benzylated ARIFLO™ or analog thereof compound has a formula,




In other embodiments, the benzylated ARIFLO™ or analog thereof compound has specific stereochemistry. Certain specific stereochemistries are shown below:



In another aspect, the present invention provides an N-benzylated
10 ROLIPRAM™ or analog thereof compound of the formula



and isolated stereoisomers, salts and solvates thereof, wherein
Bzl is a benzyl group of the formula




wherein each of the numerals 1, 2, 3, 4 and 5 may be carbon or nitrogen,
5 with the proviso that the ring is aromatic and contains at least 3 carbon atoms;

n is 5 and R¹ at each occurrence is independently selected from halogen, nitro, R², NR²_(m), and OR² wherein m = 0,1,2, or 3 and R² at each occurrence is independently selected from H, C₁-C₈alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in
10 the alkyl portion, C₃-C₇cycloalkyl, C₆-C₉polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an
15 R² group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-C₈alkoxy; and the aryl portion of an R² group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen;


A is selected from a direct bond, optionally substituted C₁-C₅alkylene, optionally substituted C₂-C₅alkenyl and optionally substituted phenylene;
20 p is 5 and R³ at each occurrence is independently selected from halogen, nitro, R⁴, NR⁴_(q), and OR⁴ wherein q = 0, 1, 2, or 3 and R⁴ at each occurrence is independently selected from H, C₁-C₈alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C₃-C₇cycloalkyl, C₆-C₉polycycloalkyl, alkylcycloalkyl, hydroxyalkyl,
25 carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R⁴ group may be optionally substituted with one or more fluorine atoms, hydroxyl or
30 C₁-C₈alkoxy; and the aryl portion of an R⁴ group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen; with the provisos that, at the meta position relative to the A group on the R³-substituted phenyl ring, R³ is alkoxy having at least three carbon atoms, excluding cyclopentyloxy, and at the para position R¹ is not hydrogen.

In a different aspect, the present invention provides an N-benzylated ROLIPRAM™ or analog thereof compound of the formula



and isolated stereoisomers, salts and solvates thereof, wherein

5 Bzl is a benzyl group of the formula



wherein each of the numerals 1, 2, 3, 4 and 5 may be carbon or nitrogen, with the proviso that the ring is aromatic and contains at least 3 carbon atoms;


10 n is 5 and R¹ at each occurrence is independently selected from halogen and OR² wherein R² at each occurrence is independently selected from H and C₁-C₈alkyl; with the provisos that the benzyl ring contain no more than three hydrogen substituents and no more than one methoxy substituent;

A is selected from a direct bond, optionally substituted C₁-C₅alkylene, optionally substituted C²-C⁵alkenyl and optionally substituted phenylene;


15 p is 5 and R³ at each occurrence is independently selected from halogen, nitro, R⁴, NR⁴_(q), and OR⁴ wherein q = 0,1,2, or 3 and R⁴ at each occurrence is independently selected from H, C₁-C₈alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C₃-C₇cycloalkyl, C₆-C₉polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, 20 carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R⁴ group may be optionally substituted with one or more fluorine atoms, hydroxyl or 25 C₁-C₈alkoxy; and the aryl portion of an R⁴ group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen.

78


In a further embodiment, the PDE4 inhibitor is an N-benzylated PDE4 inhibitor having the formula



In another aspect, the present invention provides an N-benzylated WAY-
5 PDA-641 or analog thereof compound of the formula:




and isolated stereoisomers, salts and solvates thereof, wherein Bzl is a benzyl group; and $(R^3)_p$ is as defined herein. Optionally, p is 1, 2 or 3; and in at least one occurrence, R^3 is OR^4 wherein R^4 is selected from C_1 - C_8 alkyl, C_3 - C_7 cycloalkyl, and
10 C_6 - C_9 polycycloalkyl. In a further embodiment, the PDE4 inhibitor is an N-benzylated PDE4 inhibitor having the formula




In another aspect, the present invention provides an N-benzylated Ro-20-
1724 or analog thereof compound of the formulae:

79




and isolated stereoisomers, salts and solvates thereof, wherein Bzl is a benzyl group; p is 5; and R^3 at each occurrence is independently selected from halogen, nitro, R^4 , $NR^4_{(q)}$, and OR^4 wherein $q = 0, 1, 2$, or 3 and R^4 at each occurrence is independently selected from H, C_1 - C_8 alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C_3 - C_7 cycloalkyl, C_6 - C_9 polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C_1 - C_4 alkyl and indanyl; wherein the alkyl portion of an R^4 group may be optionally substituted with one or more fluorine atoms, hydroxyl or C_1 - C_8 alkoxy; and the aryl portion of an R^4 group may be optionally substituted with C_1 - C_8 alkyl, C_1 - C_8 alkoxy or halogen. In a preferred embodiment, relative to the methylene bridge to the heterocyclic ring, R^3 at the para position is alkoxy, and R^3 at a meta position is alkoxy having at least three carbon atoms. Optionally, p is 1, 2 or 3; and in at least one occurrence, R^3 is OR^4 wherein R^4 is selected from C_1 - C_8 alkyl, C_3 - C_7 cycloalkyl, and C_5 - C_9 bicycloalkyl. In another embodiment the invention provides an N-benzylated Ro-20-1724 or analog thereof compound having the formula




20


while in another embodiment the invention provides an N-benzylated Ro-20-1724 or analog thereof compound having the formula




while in another embodiment the invention provides an N-benzylated Ro-20-1724 or analog thereof compound having the formula




- 5 while in another embodiment the invention provides an N-benzylated Ro-20-1724 or analog thereof compound having the formula




In another aspect, the present invention provides an N-benzylated RP 73401 or analog thereof compound of the formula:



and isolated stereoisomers, salts and solvates thereof, wherein Bzl is a benzyl group; and $(R^3)_p$ is as defined herein. Optionally, p is 1, 2 or 3; and in at least one occurrence, R^3 is OR^4 wherein R^4 is selected from C_1 - C_8 alkyl, C_3 - C_7 cycloalkyl, and
 5 C_6 - C_9 polycycloalkyl. In one embodiment, the N-benzylated RP 73401 or analog thereof compound has the formula:



In another aspect, the present invention provides an N-benzylated CP-80,633-A or analog thereof compound of the formulae:




10


and isolated stereoisomers, salts and solvates thereof, wherein Bzl is a benzyl group; p is 5; and R^3 at each occurrence is independently selected from nitro, R^4 , $NR^4_{(q)}$, and OR^4

wherein $q = 0, 1, 2$, or 3 and R^4 at each occurrence is independently selected from H, C_1 - C_8 alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C_3 - C_7 cycloalkyl, C_6 - C_9 polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C_1 - C_4 alkyl and indanyl; wherein the alkyl portion of an R^4 group may be optionally substituted with one or more fluorine atoms, hydroxyl or C_1 - C_8 alkoxy; and the aryl portion of an R^4 group may be optionally substituted with C_1 - C_8 alkyl, C_1 - C_8 alkoxy or halogen. In a preferred embodiment, not more than 4 of the R^3 groups are hydrogen. Optionally, p is 1, 2 or 3; and in at least one occurrence, R^3 is OR^4 wherein R^4 is selected from C_1 - C_8 alkyl, C_3 - C_7 cycloalkyl, and C_6 - C_9 polycycloalkyl. In one embodiment, the N-benzylated


15 In one embodiment the N-benzylated CP-80,633-A or analog thereof compound has the formula:




In another aspect, the present invention provides an N-benzylated AROFYLLINE™ or analog thereof of the formula:




and isolated stereoisomers, salts and solvates thereof, wherein Bzl is a benzyl group; and $(R^3)^p$ is as defined above. Optionally, p is 1, 2 or 3; and in at least one occurrence, R^3 is halogen or OR^4 wherein R^4 is selected from C_1 - C_8 alkyl, C_3 - C_7 cycloalkyl, and C_6 - C_9 polycycloalkyl. In another embodiment, the N-benzylated AROFYLLINE™ or
 5 analog thereof has the formula



while in another embodiment the N-benzylated AROFYLLINE or analog thereof has the formula



10 In another aspect, the present invention provides an N-benzylated CIPAMFYLLINE™ or analog thereof compound of the formula



and isolated stereoisomers, salts and solvates thereof, wherein Bzl is a benzyl group.

Pharmaceutical Compositions

In another embodiment, the present invention provides pharmaceutical compositions containing a benzylated compound of the invention, *i.e.*, a C- or N-benzylated compound as described herein and/or made according to the method of the present invention, in combination with a pharmaceutically-acceptable carrier, diluent or excipient. These compositions may be used for the treatment of inflammation or other conditions as disclosed herein. These compositions may also be formed into a medicament, which may be used in the treatment of, for example, inflammation.

In addition, these compositions are useful as, for example, assay standards, convenient means of making bulk shipments, or pharmaceutical compositions. An assayable amount of a compound of the invention is an amount which is readily measurable by standard assay procedures and techniques as are well known and appreciated by those skilled in the art. Assayable amounts of a compound of the invention will generally vary from about 0.001 wt% to about 98 wt% of the entire weight of the composition. Inert carriers include any material which does not degrade or otherwise covalently react with a compound of the present invention. Examples of suitable inert carriers are water; aqueous buffers, such as those which are generally useful in High Performance Liquid Chromatography (HPLC) analysis; organic solvents, such as acetonitrile, ethyl acetate, hexane and the like; and pharmaceutically acceptable carriers.

"Pharmaceutically acceptable carriers" for therapeutic use are well known in the pharmaceutical art, and are described, for example, in Remington's Pharmaceutical Sciences, Mack Publishing Co. (A.R. Gennaro edit. 1985). For example, sterile saline and phosphate-buffered saline at physiological pH may be used. Preservatives, stabilizers, dyes and even flavoring agents may be provided in the pharmaceutical composition. For example, sodium benzoate, sorbic acid and esters of p-hydroxybenzoic acid may be added as preservatives. Id. at 1449. In addition, antioxidants and suspending agents may be used. Id.

Thus, the present invention provides a pharmaceutical or veterinary composition (hereinafter, simply referred to as a pharmaceutical composition) containing a benzylated compound of the invention as described above, in admixture with a pharmaceutically acceptable carrier. The invention further provides a composition, preferably a pharmaceutical composition, containing an effective amount

of a compound of the present invention, as described above, in association with a pharmaceutically acceptable carrier.

The pharmaceutical compositions of the present invention may be in any form which allows for the composition to be administered to a patient. For example, the composition may be in the form of a solid, liquid or gas (aerosol). Typical routes of administration include, without limitation, oral, topical, parenteral, sublingual, rectal, vaginal, intranasal, and inhalation. The term parenteral as used herein includes subcutaneous injections, intravenous, intramuscular, intrasternal injection or infusion techniques. Pharmaceutical composition of the invention are formulated so as to allow the active ingredients contained therein to be bioavailable upon administration of the composition to a patient. Compositions that will be administered to a patient take the form of one or more dosage units, where for example, a tablet may be a single dosage unit, and a container of a compound of the present invention in aerosol form may hold a plurality of dosage units.

Materials used in preparing the pharmaceutical compositions should be pharmaceutically pure and non-toxic in the amounts used. It will be evident to those of ordinary skill in the art that the optimal dosage of the active ingredient(s) in the pharmaceutical composition will depend on a variety of factors. Relevant factors include, without limitation, the type of subject (*e.g.*, human), the particular form of the active ingredient, the manner of administration and the composition employed.

In general, the pharmaceutical composition includes a benzylated compound as described herein, in admixture with one or more carriers. The carrier(s) may be particulate, so that the compositions are, for example, in tablet or powder form. The carrier(s) may be liquid, with the compositions being, for example, an oral syrup or injectable liquid. In addition, the carrier(s) may be gaseous, so as to provide an aerosol composition useful in, *e.g.*, inhalatory administration.

When intended for oral administration, the composition is preferably in either solid or liquid form, where semi-solid, semi-liquid, suspension and gel forms are included within the forms considered herein as either solid or liquid.

As a solid composition for oral administration, the composition may be formulated into a powder, granule, compressed tablet, pill, capsule, chewing gum, wafer or the like form. Such a solid composition will typically contain one or more inert diluents or edible carriers. In addition, one or more of the following adjuvants may be present: binders such as carboxymethylcellulose, ethyl cellulose, microcrystalline cellulose, or gelatin; excipients such as starch, lactose or dextrins, disintegrating agents such as alginic acid, sodium alginate, Primogel, corn starch and

the like; lubricants such as magnesium stearate or Sterotex; glidants such as colloidal silicon dioxide; sweetening agents such as sucrose or saccharin, a flavoring agent such as peppermint, methyl salicylate or orange flavoring, and a coloring agent.

When the composition is in the form of a capsule, *e.g.*, a gelatin capsule,
5 it may contain, in addition to materials of the above type, a liquid carrier such as polyethylene glycol, cyclodextrin or a fatty oil.

The composition may be in the form of a liquid, *e.g.*, an elixir, syrup, solution, emulsion or suspension. The liquid may be for oral administration or for delivery by injection, as two examples. When intended for oral administration,
10 preferred composition contain, in addition to the present compounds, one or more of a sweetening agent, preservatives, dye/colorant and flavor enhancer. In a composition intended to be administered by injection, one or more of a surfactant, preservative, wetting agent, dispersing agent, suspending agent, buffer, stabilizer and isotonic agent may be included.

15 The liquid pharmaceutical compositions of the invention, whether they be solutions, suspensions or other like form, may include one or more of the following adjuvants: sterile diluents such as water for injection, saline solution, preferably physiological saline, Ringer's solution, isotonic sodium chloride, fixed oils such as synthetic mono or diglycerides which may serve as the solvent or suspending medium,
20 polyethylene glycols, glycerin, cyclodextrin, propylene glycol or other solvents; antibacterial agents such as benzyl alcohol or methyl paraben; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid; buffers such as acetates, citrates or phosphates and agents for the adjustment of tonicity such as sodium chloride or dextrose. The parenteral preparation can be
25 enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic. Physiological saline is a preferred adjuvant. An injectable pharmaceutical composition is preferably sterile.

A liquid composition intended for either parenteral or oral administration should contain an amount of a compound of the present invention such that a suitable
30 dosage will be obtained. Typically, this amount is at least 0.01% of a compound of the invention in the composition. When intended for oral administration, this amount may be varied to be between 0.001% and about 97% of the weight of the composition. Preferred oral compositions contain between about 0.1% and about 50% of the active compound of the present invention. Preferred compositions and preparations according
35 to the present invention are prepared so that a parenteral dosage unit contains between 0.1% to 5% by weight of active compound.

The pharmaceutical composition may be intended for topical administration, in which case the carrier may suitably comprise a solution, emulsion, ointment or gel base. The base, for example, may comprise one or more of the following: petrolatum, lanolin, polyethylene glycols, beeswax, mineral oil, diluents
5 such as water and alcohol, and emulsifiers and stabilizers. Thickening agents may be present in a pharmaceutical composition for topical administration. If intended for transdermal administration, the composition may include a transdermal patch or iontophoresis device. Topical formulations may contain a concentration of a benzylated compound of the present invention of from about 0.01% to about 10% w/v (weight per
10 unit volume).

The composition may be intended for rectal administration, in the form, *e.g.*, of a suppository which will melt in the rectum and release the drug. The composition for rectal administration may contain an oleaginous base as a suitable nonirritating excipient. Such bases include, without limitation, lanolin, cocoa butter
15 and polyethylene glycol.

The composition may include various materials which modify the physical form of a solid or liquid dosage unit. For example, the composition may include materials that form a coating shell around the active ingredients. The materials which form the coating shell are typically inert, and may be selected from, for example,
20 sugar, shellac, and other enteric coating agents. Alternatively, the active ingredients may be encased in a gelatin capsule.

The composition in solid or liquid form may include an agent which binds to the active component(s) and thereby assists in the delivery of the active components. Suitable agents which may act in this capacity include a monoclonal or
25 polyclonal antibody, a protein or a liposome.

The pharmaceutical composition of the present invention may consist of gaseous dosage units, *e.g.*, it may be in the form of an aerosol. The term aerosol is used to denote a variety of systems ranging from those of colloidal nature to systems consisting of pressurized packages. Delivery may be by a liquefied or compressed gas or
30 by a suitable pump system which dispenses the active ingredients. Aerosols of compounds of the invention may be delivered in single phase, bi-phasic, or tri-phasic systems in order to deliver the active ingredient(s). Delivery of the aerosol includes the necessary container, activators, valves, subcontainers, spacers and the like, which together may form a kit. Preferred aerosols may be determined by one skilled in the art,
35 without undue experimentation.

Whether in solid, liquid or gaseous form, the pharmaceutical composition of the present invention may contain one or more known pharmacological agents used in the treatment of inflammation (including arthritis).

The pharmaceutical compositions may be prepared by methodology well known in the pharmaceutical art.

A composition intended to be administered by injection can be prepared by combining a benzylated compound as described herein with water so as to form a solution. A surfactant may be added to facilitate the formation of a homogeneous solution or suspension. Surfactants are compounds that non-covalently interact with the benzylated compounds of the invention so as to facilitate dissolution or homogeneous suspension of the active compound in the aqueous delivery system.

In a preferred embodiment, a benzylated PDE4 inhibitor according to the present invention, or a benzylated PDE4 inhibitor prepared according to the present invention, has an IC_{50} of 600 nM (nanomolar) or less, more preferably 400 nM or less.

Methods for Assessment of PDE4 Activity, HARBS Activity and Emetogenic Potential

Various *in vitro* tests, which are well known to one of ordinary skill in the art, may be used to determine whether a particular compound is a PDE inhibitor. Some tests are specific to a particular PDE isozyme or conformer, while other tests are less specific. A few of these tests are described below. The identification of other testing protocols is readily achieved by reviewing the scientific and patent literature, including Burnouf, C. et al. "Phosphodiesterase 4 Inhibitors" *Annual Reports in Medicinal Chemistry*, Vol. 33, Chap. 10, pp 91-109, 1998 (Bristol, J.A., ed.); Essayan, D. M. "Cyclic Nucleotide Phosphodiesterase (PDE) Inhibitors and Immunomodulation" *Biochemical Pharmacology* 57:965-973, 1999; Souness, J. E. and Foster, M. "Potential of phosphodiesterase type IV inhibitors in the treatment of rheumatoid arthritis" *Idrugs* 1(5):541-553, 1998; Souness, J.E. et al. "Immunosuppressive and anti-inflammatory effect of cAMP phosphodiesterase (PDE) type 4 inhibitors" *Immunopharmacology* 47: 127-162, 2000; and Torphy, T. J. "Phosphodiesterase Isozymes" *Am J. Respir. Crit. Care Med.* 157:351-370, 1998, as well as the numerous references cited in these articles.

A test compound may be screened for activity against 11 of the major classes of mammalian cyclic nucleotide phosphodiesterase (termed PDE 1 through 11). These PDE's may use cAMP or cGMP or both cyclic nucleotides as substrates. The broad specificity PDE inhibitor 3-isobutyl-1-methylxanthine (IBMX; Sigma; Catalogue No. 17018) or like compound may be used as a positive control in most assays. PDE's

for some of the various assays may be fully or partially purified from the following cells/tissues: PDE 1 (bovine heart), PDE 2 (human platelets), PDE 3 (human platelets), PDE4 (human promonocytic U937 cells) and PDE 5 (human platelets).

In the following descriptions, amounts are exemplary, and may be varied. U937 cytoplasmic extracts may be prepared by sonicating U937 cells (ATCC: Catalogue No. CRL-159) in lysis buffer (M-PER mammalian protein extraction reagent, Pierce catalog # 78501 containing 10% Protease Inhibitor Cocktail For Use with Mammalian Cell and Tissue Extracts, Sigma catalog # P8340). Sonicated cell extracts are then centrifuged at 70,000 g for 30 minutes and supernatants removed. Sucrose is added to a final concentration of 0.25 M, then the extracts are aliquoted and stored at -80°C.

PDE reactions are performed using the Phosphodiesterase [^3H] cAMP SPA enzyme assay as described by the manufacturer (Amersham Pharmacia Biotech; <http://www.apbiotech.com>; catalog # 7090). Briefly, the assay is run for 60 minutes at room temperature in 50 μL volumes in 1 μM [^3H] cAMP, 50 mM Tris Cl, 10 mM MgCl_2 pH 7.5. U937 extract is added such that less than 10% of substrate was consumed. Test compound or vehicle is added to the desired concentration. Typically, compounds are tested at six 10-fold dilutions ranging from 100 μM to 1 nM. Reactions are performed in duplicate. Reactions are terminated with 50 μl of Yttrium SPA beads and then counted on a Packard Scintillation counter (TopCount™) for 1 minute.

Utilizing the PDE assay conditions described above with PDE4, typical PDE4 inhibitors such as ROLIPRAM™ and Ro-20-1724 (Calbiochem: Catalogue No. 557502) give IC_{50} values in agreement with those found in the literature (reviewed in Schudt et al., 1996).

Inhibition of PDE4 (or more accurately, specific isoforms of PDE4) with subsequent elevation of intracellular cAMP and protein kinase A activation is a therapeutic target in inflammatory or autoimmune diseases where the causal cells or tissues involved predominantly express this PDE isoform. With respect to rheumatoid arthritis, the PDE4 inhibitor ROLIPRAM™ has been shown to be active in animal models of the disease such as collagen-induced arthritis in the rat (Nyman et al., *Clin. Exp. Immunol.* 108(3), 415-419, 1997).

Compounds including a benzyl group according to the process of the present invention typically demonstrate PDE4 inhibition with reduced emetogenic potential, relative to the corresponding non benzylated analog. The efficacy of a compound as a PDE4 inhibitor with diminished emetogenic potential may be measured by determining the extent to which the compound displaces ROLIPRAM™ from its preferred

binding site (known as the High Affinity Binding Site, or HARBs) on PDE4. Animal models have shown that a compound's antiinflammatory efficacy as a PDE4 inhibitor is highly correlated with its potency at inhibiting PDE4 catalytic activity rather than its ability to displace [³H]-ROLIPRAM™ from a high affinity binding site from cells within
5 the brain and central nervous system (Duplantier 1996, Barnette 1996). Compounds having benzyl groups according to the present invention displayed a low affinity for the HARBS conformer of PDE4 suggesting that these compounds are not likely to be plagued by mechanism-associated side-effects associated with first generation PDE4 inhibitors such as ROLIPRAM™.

10 In order to perform this assay, female CD1 mice are sacrificed via the intraperitoneal injection of 100 µL ethanol, and the brain tissue is homogenized in 5 mL of ice-cold Tris-HCl, pH 8.00 supplemented with 1.2 mM MgCl₂, 1 mM benzamidine (Sigma; Catalogue No. B 6506) and 0.1 mM PMSF (Sigma; Catalogue No. P 7626). The suspension is centrifuged twice at 30,000 × G at 4°C and then the supernatant discarded.
15 The pellet is resuspended in buffer, and adjusted to a protein concentration of 0.5 mg/mL. Compounds to be tested are dissolved in DMSO and pipetted in triplicate into a 96 well microplate at concentrations ranging from 1 to 30,000 nM. 10 mL of membrane preparation is supplemented with 100 µL of 0.235 µM [³H]-ROLIPRAM™ in DMSO, and 100 µL is dispensed into each well of the microplate. The plate is incubated at 4°C
20 for 1 hour. Contents of the plate are aspirated through a Whatman GF/C filterplate, and rinsed with 4×200 µL ice-cold buffer. The plate is dried overnight, and then 30 µL of Microscint 20 (Packard; Catalogue No. 6013621) is added to each well, and plate is read in the scintillation counter with a sampling time of 2 minutes/well. Values representing non-specific binding (defined by counts obtained using 20 µM ROLIPRAM™) are
25 subtracted from all data points. Triplicate determinations should be performed at each concentration. The PDE4:HARBS value indicates the ratio of the IC₅₀ concentration required to inhibit catalytic activity to the concentration required to displace 50% of ROLIPRAM™ from the high affinity binding site.

Under these assay conditions ROLIPRAM™ is able to displace ³H-
30 ROLIPRAM™ from a high-affinity binding site in mouse brain with an IC₅₀ of about 6 nM (data not shown). Thus, ROLIPRAM™ binds with 65 fold greater affinity to its high affinity site than the concentration required for half-maximal inhibition of PDE4 catalytic activity. This preferential affinity for HARBS over the catalytic conformer has been correlated with the negative side effects of first generation PDE4 inhibitors;
35 namely emesis and CNS effects.

Benzylated compounds according to the present invention are less potent at binding to this HARBS site than ROLIPRAM™. For example, see the LPDE4:HPDE4 ratio reported for compound 25 in Table 1 in the Example below. Thus, compounds of the present invention may display *in-vivo* emetogenic effects that are much less than ROLIPRAM™, Ro 20-1724 or other first generation PDE4 inhibitors.

Utility of Compounds and Compositions of the Present Invention

The compounds disclosed herein, or compositions comprising one of more of these compounds and a pharmaceutically acceptable carrier, diluent or excipient, may be used in a method for various treatments and prophylactic effects. For instance, the compounds of the invention may be used for treating or preventing an inflammatory condition or disease in a patient, where the method comprises administering to the patient in need thereof an amount of a compound or composition according to the present invention, where the amount is effective to treat or prevent the inflammatory condition or disease of the patient. The compounds and compositions of the present invention are particularly advantageous in that they provide the desired activity, *e.g.*, treatment of inflammation, without attendant undesirable emetogenic side-effects.

The inflammatory condition or disease may be an autoimmune condition or disease; the inflammatory condition or disease may involve acute or chronic inflammation of bone and/or cartilage compartments of joints; the inflammatory condition or disease may be an arthritis selected from rheumatoid arthritis, gouty arthritis or juvenile rheumatoid arthritis; the inflammatory condition or disease may be asthma; the condition or disease may be associated with the dysregulation of T-cells; the condition or disease may be associated with elevated levels of inflammatory cytokines (*e.g.*, wherein the inflammatory cytokine is IL-2, IL-4, IL-5, or IL-12, or wherein the inflammatory cytokine is IFN- γ , or wherein the inflammatory cytokine is TNF- α); the inflammatory condition or disease may be multiple sclerosis; the inflammatory condition or disease may be pulmonary sarcoidosis; the inflammatory condition or disease may be ocular inflammation or allergy; the inflammatory condition or disease may be an inflammatory bowel disease (*e.g.*, Crohn's disease or ulcerative colitis); and the inflammatory condition or disease may be an inflammatory cutaneous disease (*e.g.*, psoriasis or dermatitis).

Furthermore, the present invention provides a method for modulating intracellular cyclic adenosine 3',5'-monophosphate levels within a patient, comprising

administering to a patient in need thereof an amount of a compound or composition according to the present invention, wherein the amount is effective to modulate the intracellular cyclic adenosine 3',5'-monophosphate levels of the patient. The patient may have an inflammatory condition or disease.

5 Furthermore, the present invention provides a method for treating or preventing a disease or condition in a patient, where the disease or condition is associated with pathological conditions that are modulated by inhibiting enzymes associated with secondary cellular messengers, the method comprising administering to
10 a patient in need thereof an amount of a compound or a composition of the present invention, wherein the amount is effective to treat or prevent a disease or condition associated with pathological conditions that are modulated by inhibiting enzymes associated with secondary cellular messengers. The enzyme may be a cyclic AMP phosphodiesterase; or the enzyme may be a phosphodiesterase 4.

Furthermore, the present invention provides a method of treating or
15 preventing transplant rejection in a patient, comprising administering to a patient in need thereof an amount of a compound or composition of the present invention, where the amount is effective to treat or prevent transplant rejection in the patient. The rejection may be due to graft versus host disease.

Furthermore, the present invention provides a method of treating or
20 preventing uncontrolled cellular proliferation in a patient, comprising administering to a patient in need thereof an amount of a compound or composition according to the present invention, where the amount is effective to treat or prevent uncontrolled cellular proliferation in the patient. The uncontrolled cellular proliferation may be caused by a cancer selected from leukemia and solid tumors.

25 Furthermore, the present invention provides a method of treating or preventing conditions associated with the central nervous system (CNS) in a patient, comprising administering to a patient in need thereof an amount of a compound or composition according to the present invention, where the amount is effective to treat or prevent conditions associated with the central nervous system (CNS) in the patient. The
30 condition associated with the central nervous system (CNS) may be depression.

In one aspect the present invention provides a method for treating or preventing an inflammatory condition or disease in a patient. The method includes administering to the patient in need thereof an amount of a benzylated PDE4 inhibitor or composition containing a benzylated PDE4 inhibitor as described in the various
35 inventive aspects and embodiments described herein, where the amount is effective to treat or prevent the inflammatory condition or disease of the patient. Optionally, the

benzylated PDE4 inhibitor inhibits an enzyme selected from phosphodiesterase 4 A, B, C or D or any combination thereof including all splice variants of PDE4 A, B, C and D. In one aspect, the inflammatory condition or disease is an autoimmune condition or disease. In another aspect, the inflammatory condition or disease involves acute or chronic inflammation of bone and/or cartilage compartments of joints. In another aspect, the inflammatory condition or disease is an arthritis selected from rheumatoid arthritis, gouty arthritis or juvenile rheumatoid arthritis. In another aspect, the inflammatory condition or disease is asthma. In another aspect, the inflammatory condition or disease is associated with the dysregulation of T-cells. In another aspect, the inflammatory condition or disease is associated with elevated levels of inflammatory cytokines. Optionally, the inflammatory cytokine is IL-2, IL-4, IL-5, or IL-12. Optionally, the inflammatory cytokine is IFN- γ . Optionally, the inflammatory cytokine is TNF- α . In another aspect, the inflammatory condition or disease is multiple sclerosis. In another aspect, the inflammatory condition or disease is pulmonary sarcoidosis. In another aspect, the inflammatory condition or disease is ocular inflammation or allergy. In another aspect, the inflammatory condition or disease is an inflammatory bowel disease. Optionally, the inflammatory bowel disease is Crohn's disease or ulcerative colitis. In another aspect, the inflammatory condition or disease is an inflammatory cutaneous disease. Optionally, the inflammatory cutaneous disease is psoriasis or dermatitis. In another aspect, the inflammatory condition or disease is chronic obstructive pulmonary disease (COPD), bronchitis, emphysema or acute respiratory distress syndrome (ARDS).

In another aspect the present invention provides a method for treating or preventing a disease or condition in a patient, where the disease or condition is associated with pathological conditions that are modulated by inhibiting enzymes associated with secondary cellular messengers. The inventive method includes administering to the patient in need thereof an amount of a benzylated PDE4 inhibitor or composition containing a benzylated PDE4 inhibitor as described in the various inventive aspects and embodiments described herein, wherein the amount is effective to treat or prevent a disease or condition associated with pathological conditions that are modulated by inhibiting enzymes associated with secondary cellular messengers. In one aspect, the enzyme is a cyclic AMP phosphodiesterase. In another aspect, the enzyme is phosphodiesterase 4.

In another aspect the present invention provides a method for treating or preventing transplant rejection in a patient. The inventive method includes administering to the patient in need thereof an amount of a benzylated PDE4 inhibitor

or composition containing a benzylated PDE4 inhibitor as described in the various inventive aspects and embodiments described herein, where the amount is effective to treat or prevent transplant rejection in the patient. In one aspect, the rejection is due to graft versus host disease.

5 In another aspect the present invention provides a method for treating or preventing uncontrolled cellular proliferation in a patient. The method includes administering to the patient in need thereof an amount of a benzylated PDE4 inhibitor or composition containing a benzylated PDE4 inhibitor as described in the various inventive aspects and embodiments described herein, where the amount is effective to
10 treat or prevent uncontrolled cellular proliferation in the patient. In one aspect, the uncontrolled cellular proliferation is caused by a cancer selected from leukemia and solid tumors.

In another aspect, the present invention provides a method of treating or preventing conditions associated with the central nervous system (CNS) in a patient.
15 The method includes administering to the patient in need thereof an amount of a benzylated PDE4 inhibitor or composition containing a benzylated PDE4 inhibitor as described in the various inventive aspects and embodiments described herein, where the amount is effective to treat or prevent conditions associated with the central nervous system (CNS) in the patient. In one aspect, the condition associated with the CNS is
20 depression. In another aspect, the condition associated with the CNS is long-term memory potentiation and learning enhancement.

In another aspect, the present invention provides a method of treating or preventing diseases in a patient, where the disease is associated with viral infection. The method includes administering to the patient in need thereof an amount of a
25 benzylated PDE4 inhibitor or composition containing a benzylated PDE4 inhibitor as described in the various inventive aspects and embodiments described herein, where the amount is effective to treat or prevent diseases associated with viral infection in the patient. In one aspect, the viral infection is due to the human immunodeficiency virus (HIV) and the disease is acquired immunodeficiency syndrome (AIDS).

30 In another aspect, the present invention provides a method of treating or preventing diseases in a patient, where the disease is associated with infection by a parasite. The method includes administering to the patient in need thereof an amount of a benzylated PDE4 inhibitor or composition containing a benzylated PDE4 inhibitor as described in the various inventive aspects and embodiments described herein, where the
35 amount is effective to treat or prevent diseases associated with infection of the patient

by a parasite. In one aspect, the parasitic infection is due to the trypanosoma brucei and the disease is African sleeping sickness disease.

In another aspect, the present invention provides a method of treating or preventing cystic fibrosis in a patient. The method includes administering to the patient in
5 need thereof an amount of a benzylated PDE4 inhibitor or composition containing a benzylated PDE4 inhibitor as described in the various inventive aspects and embodiments described herein, where the amount is effective to treat or prevent cystic fibrosis in the patient.

In a method of the present invention, a benzylated compound, or a
10 composition comprising one or more benzylated and a pharmaceutically acceptable carrier, diluent or excipient, may, although need not, achieve one or more of the following desired results in the subject to whom has been administered a benzylated compound as defined above, or a composition containing one of these compounds and a pharmaceutically acceptable carrier, diluent or excipient:

- 15 1. Inhibition of cyclic-AMP phosphodiesterase 4;
2. Low ratio of IC_{50} PDE4(cat): IC_{50} PDE4(HARBS);
3. Inhibition of TNF- α production;
4. Inhibition of edema;
5. Inhibition of reactive oxygen species generation from primary
20 neutrophils;
6. Oxygen radical scavenging;
7. Potentiate induction of CRE-mediated transcription activity in human monocytic cells;
8. Inhibition of PDE, preferably PDE4, PDE3, or PDE3 and PDE4;
- 25 9. Inhibition of cytokine production by activated T-cell subsets;
10. Inhibition of neutrophil myeloperoxidase release;
11. Inhibition of neutrophil chemotaxis;
12. Inhibition of graft rejection;
13. Inhibition of clinical and histopathological parameters of disease in
30 inflammatory bowel disease;
14. Inhibition of clinical and histopathological parameters of arthritis in a murine collagen-induced arthritis model; and
15. Inhibition of clinical and histopathological parameters of disease in an allergan-induced guinea-pig model of acute bronchoconstriction.

35 Thus, the inventive method may be used to treat inflammation, including both acute and chronic inflammation as well as certain proliferative disorders (cancers).

As used herein, inflammation includes, without limitation, ankylosing spondylitis, arthritis (where this term encompasses over 100 kinds of rheumatic diseases), asthma, Crohn's disease, fibromyalgia syndrome, gout, inflammations of the brain (including multiple sclerosis, AIDS dementia, Lyme encephalopathy, herpes encephalitis, Creutzfeld-Jakob disease, and cerebral toxoplasmosis), emphysema, inflammatory bowel disease, irritable bowel syndrome, ischemia-reperfusion injury, juvenile erythematosis, pulmonary sarcoidosis, Kawasaki disease, osteoarthritis, pelvic inflammatory disease, psoriatic arthritis (psoriasis), rheumatoid arthritis, psoriasis, tissue/organ transplant, scleroderma, spondyloarthropathies, systemic lupus erythematosis, pulmonary sarcoidosis, and ulcerative colitis. As used herein, proliferative disorders includes, without limitation, all leukemias and solid tumors that are susceptible to undergoing differentiation or apoptosis upon interruption of their cell cycle.

A method of the present invention is the administration of a therapeutically effective amount of a benzylated compound as described herein including salts, compositions etc. thereof. As used herein, the actual amount encompassed by the term "therapeutically effective amount" will depend on the route of administration, the type of warm-blooded animal being treated, and the physical characteristics of the specific warm-blooded animal under consideration. These factors and their relationship to determining this amount are well known to skilled practitioners in the medical arts. This amount and the method of administration can be tailored to achieve optimal efficacy but will depend on such factors as weight, diet, concurrent medication and other factors that those skilled in the medical arts will recognize.

An effective amount of a compound or composition of the present invention will be sufficient to treat inflammation in a warm-blooded animal, such as a human. Methods of administering effective amounts of anti-inflammatory agents are well known in the art and include the administration of inhalation, oral or parenteral forms. Such dosage forms include, but are not limited to, parenteral solutions, tablets, capsules, sustained release implants and transdermal delivery systems; or inhalation dosage systems employing dry powder inhalers or pressurized multi-dose inhalation devices.

The dosage amount and frequency are selected to create an effective level of the agent without harmful effects. It will generally range from a dosage of about 0.001 to 100 mg/Kg/day, and typically from about 0.01 to 10 mg/Kg/day where administered orally or intravenously. Also, the dosage range will be typically from about 0.001 to 1 mg/Kg/day where administered intranasally or by inhalation.

The following examples are offered by way of illustration and not by way of limitation.

Unless otherwise stated, flash chromatography and column chromatography may be accomplished using Merck silica gel 60 (230–400 mesh). Flash chromatography may be carried out according to the procedure set forth in: "Purification of Laboratory Chemicals", 3rd. edition, Butterworth-Heinemann Ltd., Oxford (1988), Eds. D. D. Perrin and W. L. F. Armarego, page 23. Column chromatography refers to the process whereby the flow rate of eluent through a packing material is determined by gravity. In all cases flash chromatography and radial chromatography may be used interchangeably. Radial chromatography is performed using silica gel on a Chromatotron Model # 7924T (Harrison Research, Palo Alto, California). Unless otherwise stated, quoted R_f values are obtained by thin layer chromatography using Silica Gel 60 F₂₅₄ (Merck KGaA, 64271, Darmstadt, Germany).

Also, unless otherwise stated, chemical reactants and reagents were obtained from standard chemical supply houses, such as Aldrich (Milwaukee, WI; www.aldrich.sial.com); EM Industries, Inc. (Hawthorne, NY; www.emscience.com); Fisher Scientific Co. (Hampton, NH; www.fischer1.com); and Lancaster Synthesis, Inc. (Windham, NH; www.lancaster.co.uk). Gases were obtained from Praxair (Vancouver, B.C.). Cell lines, unless otherwise stated, were obtained from public or commercial sources, e.g., American Tissue Culture Collection (ATCC, Rockville, MD).


EXAMPLES

EXAMPLE 1

FORMATION OF BENZYLATION AGENT FROM CORRESPONDING BENZYL ALCOHOL

25

Scheme 2



Synthesis of Compound 2

To a rapidly stirred slurry of 3-methoxy-4-hydroxybenzyl alcohol 1 (30.0 g, 195 mmol), potassium carbonate (62.2 g, 450 mmol), and 18-crown-6 (0.40 g, 1 mol%) in toluene (350 mL) was added a solution of benzyl bromide (BnBr, 25.6 g, 150 mmol) in toluene (150 mL) over 20 min. The reaction mixture was refluxed for 16 hours, after which the mixture was diluted with diethyl ether (400 mL) and washed successively with NaOH (1 N, 2 x 250 mL), saturated aqueous NaHCO₃ (2 x 250 mL), and brine (2 x 300 mL). The diethyl ether layer was dried over anhydrous MgSO₄, and the solvent was removed to provide a pale yellow solid (42.1 g) which was crystallized with EtOAc and hexanes to give compound 2 (32.7 g, 77%) as a white crystalline solid.

Synthesis of Compound 3


To a solution of alcohol 2 (3.20 g, 13.1 mmol) in anhydrous diethyl ether (15 mL) was added PBr₃ (1.77 g, 6.55 mmol) in one portion, and the resulting mixture was stirred at room temperature for 3 hours. The mixture was diluted with diethyl ether (40 mL) and washed with H₂O (2 x 30 mL), saturated NaHCO₃ (2 x 30 mL), and brine (2 x 30 mL). The ether layer was dried over anhydrous MgSO₄, and the solvent was removed under reduced pressure to afford compound 3 (4.02 g, 100%) as a light yellow solid.

EXAMPLE 2

20 FORMATION OF BENZYLATION AGENT FROM CORRESPONDING BENZYL ALDEHYDE

Scheme 3 illustrates several procedures for converting commercially available benzyl aldehydes into the corresponding benzyl alcohol. Suitable solvents may be employed in any of these reactions.

99

Scheme 3

Synthesis of Compound 14

To a suspension of 4-hydroxy-3-nitrobenzaldehyde 13 (3.00 g, 17.95 mmol), potassium carbonate (3.73 g, 26.93 mmol) in DMF (300 mL) was slowly added benzyl bromide (2.85 mL, 23.96 mmol). The reaction mixture was stirred at 65°C for 18 hours. After cooling to room temperature, the mixture was diluted with water (140 mL) and extracted with diethyl ether (3 x 150 mL). The combined organic layers were washed with water (150 mL) and brine (150 mL). After drying over anhydrous MgSO₄, filtration and evaporation of the filtrate *in vacuo* gave crude compound 14 (4.39 g, 95%) which was used for the next reaction without further purification.

10 Synthesis of Compound 15

Compound 14 (4.30 g, 16.72 mmol) was dissolved in EtOH/CH₂Cl₂ (1:1, 50 mL) and cooled to 0°C. NaBH₄ (0.63 g, 16.72 mmol) was added portionwise. After the addition was completed, the ice-water bath was removed and the reaction mixture was stirred at room temperature for 2 hours. Water (40 mL) was added and the mixture was extracted with CH₂Cl₂ (3 x 60 mL). The combined organic layers were washed with brine (50 mL) and dried over anhydrous MgSO₄. Removal of the solvent gave a pale yellow solid which was purified by silica gel column chromatography (hexanes/EtOAc, 1:1) to give compound 15 (4.31 g, 99%) as a pale yellow solid.

Synthesis of Compound 16


20 To a solution of compound 15 (4.30 g, 16.59 mmol) in anhydrous diethyl ether (40 mL) was slowly added PBr₃ (0.79 mL, 8.30 mmol) *via* syringe, and the resulting mixture was stirred at room temperature for 3 hours. The mixture was diluted with EtOAc (100 mL) and washed with saturated aqueous NaHCO₃ (2 x 50 mL) and brine (2 x 50 mL). The organic layer was dried over anhydrous MgSO₄, and the solvent was removed under reduced pressure to afford compound 16 (5.07 g, 95%) as a pale yellow solid.

EXAMPLE 3

FORMATION OF BENZYLATED PDE4 INHIBITOR FROM BENZYLATION AGENT AND
CORRESPONDING NON-BENZYLATED INHIBITOR

30 A. Benzylation using various halide compounds to provide the desired benzylated products may be achieved as shown in Scheme 4. In Scheme 4, compound 17 is benzylated with commercially available 3,4-difluorobenzyl bromide to give the desired compound 18 in 59% yield.

101

Scheme 4Synthesis of Compound 18


5 *n*-Butyllithium (2.5 M solution in hexanes, 0.56 mL, 1.40 mmol) was added to a solution of diisopropylamine (0.20 mL) in dry THF (10 mL) at -78°C . The mixture was stirred at -78°C for 1 hour, then HMPA (0.33 mL, 1.91 mmol) was added, followed by addition of a solution of compound 17 (0.30 g, 1.27 mmol) in THF (3 mL). After 1 hour, a solution of 3,4-difluoro benzyl bromide (purchased from Aldrich

10 Chemical Company, Inc., 0.32 mL, 2.54 mmol) was added in one portion to the reaction, and the resulting mixture was stirred at -78°C for an additional 4 hours. The excess base was quenched with saturated aqueous NH_4Cl (10 mL), and the resulting solution was extracted with EtOAc (3 x 20 mL). The combined organic layer was washed with saturated NaCl (2 x 30 mL), dried over MgSO_4 , filtered and the filtrate

15 evaporated to dryness. The residue was purified by column chromatography on silica gel (hexanes/EtOAc, 2:1) to give compound 18 (0.27 g, 59%) as a white solid.

B. In Scheme 5, compound 17 is benzylated with 3,4-dibenzyloxy benzyl bromide (prepared by treatment the corresponding alcohol with PBr_3), followed by hydrogenation using 10% Pd/C as catalyst to give the desired compound 21 in good

20 yield.

Scheme 5

Synthesis of Compound 19

To a solution of 3,4-dibenzyloxybenzyl alcohol (1.35 g, 4.21 mmol) in anhydrous diethyl ether (25 mL) was added PBr_3 (0.20 mL, 2.11 mmol) in one portion, and the resulting mixture was stirred at room temperature for 3 hours. The mixture was
5 diluted with diethyl ether (50 mL) and washed with H_2O (2 x 30 mL), saturated NaHCO_3 (2 x 30 mL), and brine (2 x 30 mL). The ether layer was dried over anhydrous MgSO_4 , and the solvent was removed under reduced pressure to afford compound 19 (1.47 g, 91%) as a light yellow oil.

Synthesis of Compound 20

10 *n*-Butyllithium (2.5 M solution in hexanes, 0.38 mL, 0.931 mmol) was added to a solution of diisopropylamine (0.14 mL 0.999 mmol) in dry THF (3 mL) at -78°C . The mixture was stirred at -78°C for 1 hour, then HMPA (0.22 mL, 1.27 mmol) was added, followed by adding a solution of compound 17 (200.0 mg, 0.846 mmol) in THF (3 mL). After 1 hour, a solution of 3,4-dibenzyloxy benzyl bromide (compound
15 19, 248.5 mg, 0.80 mmol) in THF (1 mL) was added in one portion to the reaction, and the resulting mixture was stirred at -78°C for an additional 4 hours. The excess base was quenched with saturated aqueous NH_4Cl (10 mL), and the resulting solution was extracted with EtOAc (3 x 20 mL). The combined organic layer was washed with saturated NaCl (2 x 30 mL), dried over MgSO_4 , filtered and the filtrate evaporated to
20 dryness. The residue was purified by column chromatography on silica gel (hexanes/EtOAc, 2:1) to give compound 20 (0.31 g, 67%) as a colorless oil.

Synthesis of Compound 21

A mixture of compound 20 (0.20 g, 0.37 mmol) and 10% Pd/C (25 mg) in EtOAc/AcOH (4:1, 5 mL) was stirred under H_2 (balloon) for 2 hours. The mixture
25 was then filtered through a celite plug and the filtrate was evaporated to dryness. The residue was purified by column chromatography on silica gel (hexanes/EtOAc, 3:2) to give compound 21 (0.093 mg, 70%) as a colorless syrup.

EXAMPLE 4


FORMATION OF ALPHA-BENZYLATED ROLIPRAMTM

30 Further exemplary synthetic methodology to provide a benzylated ROLIPRAMTM compound is described below. For example, as illustrated in Scheme 6, *N*-protection of 22 (prepared according to the literature procedure: *Journal of Medicinal Chemistry*, 1993, 36, 3274-3277) with di-*tert*-butyl dicarbonate and triethylamine in

dichloromethane provides *N*-*t*-butoxycarbonylamide derivative 23. Alkylation of compound 23 with 4-(benzyloxy)-3-methoxybenzyl bromide 3 affords compound 24. Any number of alkylations can be carried out using different halides and the appropriate base such as LDA. Removal of the *N*-BOC protecting group in with trifluoroacetic acid
5 in dichloromethane gives compound 25. Hydrogenolysis of compound 25 using 10% Pd/C as catalyst provides the desired product 26.

104

Scheme 6



Preparation of Compound 23

Di-*tert*-butyl dicarbonate (0.793 mg, 3.63 mmol) was added to a solution of compound 22 (0.50 g, 1.82 mmol), Et₃N (0.50 mL, 3.63 mmol) and DMAP (0.044 g) in CH₂Cl₂ (14 mL). The mixture was stirred at room temperature for 4 hours. The mixture was concentrated, and the residue was purified by column chromatography on silica gel eluted with hexane/EtOAc (3:2) to afford compound 23 (0.655 g, 96%) as light yellow wax.

Preparation of Compound 24

To a solution of compound 23 (0.649 g, 1.73 mmol) in dry THF (13 mL) under argon was slowly added LDA [1.20 mmol, prepared from *n*-BuLi (0.83 mL, 2.5 M solution in hexane, 1.20 mmol) and diisopropylamine (0.30 mL, 2.16 mmol)] in THF (3 mL) at -78°C. The mixture was stirred at -78°C for one hour, and then DMPU (0.60 mL, 1.50 mmol) was added to the above mixture via syringe. After 15 minutes, 4-(benzyloxy)-3-methoxybenzyl bromide 3 (1.06 g, 3.46 mmol) in THF (1.5 mL) was added. The resulting mixture was stirred for an additional 4 hours at -78°C. The excess base was quenched at 0°C with aqueous saturated NH₄Cl (20 mL), and the resulting solution was extracted with EtOAc (3 x 30 mL). The combined organic layer was washed with saturated brine (2 x 30 mL), dried over MgSO₄, filtered and the filtrate was evaporated to dryness. The residue was purified by column chromatography on silica gel eluted with hexanes/EtOAc (7:3) to give compound 24 (0.829 g, 80%) as a white foam.

Preparation of Compound 25

Trifluoroacetic acid (14 mL) was added to a solution of compound 24 (0.820 g, 1.36 mmol) in CH₂Cl₂ (14 mL). The mixture was stirred at room temperature for 2 hours, diluted with CH₂Cl₂ (100 mL) and washed with saturated NaHCO₃ solution (3 x 20 mL). The organic layer was dried over MgSO₄, filtered and the filtrate was evaporated to dryness. The residue was purified by column chromatography over silica gel using ethyl acetate as eluent to afford compound 25 (0.510 g, 75%) as white solid.

Preparation of Compound 26

A mixture of compound 25 (0.356 g, 0.710 mmol) and 10% Pd/C (36 mg) in EtOAc/AcOH (1:1, 12 mL) was stirred under H₂ (balloon pressure) for 18 hours. The mixture was filtered on celite and the filtrate was evaporated to dryness. The


residue was purified by column chromatography on silica gel eluted with Hexanes/EtOAc (10:90) to afford compound 26 (0.20 g, 84%) as a white foam.

EXAMPLE 5

FORMATION OF *N*-BENZYLATED ROLIPRAM™

- 5 An example of *N*-benzylation using benzyl halide to provide the desired product 27 is illustrated in Scheme 7. In scheme 7, compound 22 is benzylated with 4-(cyclopentyloxy)-3-methoxybenzyl bromide 44 to provide desire compound 27.

Scheme 7



10


EXAMPLE 6

FORMATION OF BETA-BENZYLATED ARIFLO™ (33)

- As illustrated in Scheme 8, compound 28 (prepared according to the literature procedure: *Journal of Medicinal Chemistry*, 1998, 41, 821-835) is alkylated with 4-(benzyloxy)-3-methoxybenzyl bromide 3 to provide compound 29, which is saponified and decarboxylated to cyclohexanone 30 with sodium chloride in hot aqueous DMSO. Cyclohexanone 30 is homologated by a Peterson-type reaction with excess 2-lithio-2-(trimethylsilyl)-1,3-dithiane (prepared from 2-TMF-1,3-dithiane and *n*-butyl lithium in THF) to provide ketene dithioacetal 31. Methanolysis of compound 31 with mercuric chloride and perchloric acid provides ester 32. Saponification of the ester 32 with potassium hydroxide in a THF/methanol/water solvent mixture, followed by acidification with aqueous HCl, is used to provide the desired product 33.

107

Scheme 8



EXAMPLE 7


FORMATION OF GAMMA-BENZYLATED ARIFLO™ (39)

- Exemplary synthetic methodology to introduce a benzyl or substituted benzyl group at the carbon γ to the carbonyl group of a compound of formula (28) is described below. For example, as illustrated in Scheme 9, compound 28 (prepared according to the literature procedure: *Journal of Medicinal Chemistry*, 1998, 41, 821-835) may be saponified and decarboxylated to compound 34 with sodium chloride in hot aqueous DMSO. The cyclohexenone ring may be generated using isopropenyl acetate and *p*-toluenesulphonic acid (TsOH) followed by Pd(OAc)₂, Bu₃Sn(OMe) and CH₂CHOCO₂CH₃. Addition of a benzyl group at the beta position of the cyclohexenone ring may be achieved by a Michael-type addition using commercially available benzylmagnesium chloride as shown in Scheme 9 to yield compound 36. Any number of substituted organomagnesium or organolithium benzyl groups may be prepared and used in Michael-type reactions according to the methods described in

Comprehensive Organic Transformations, A Guide to Functional Group Preparations, Second Edition, Richard C. Larock, John Wiley and Sons, Inc., 1999, or referenced therein. Cyclohexanone 36 is homologated by a Peterson-type reaction with excess 2-lithio-2-(trimethylsilyl)-1,3-dithiane to provide ketene dithioacetal 37. Methanolysis of
5 compound 37 with mercuric chloride and perchloric acid provides ester 38. Hydrolysis of the ester 38 with potassium hydroxide in a THF/methanol/water solvent mixture, followed by acidification with aqueous HCl, may provide the desired product 39.

109

Scheme 9




EXAMPLE 8

FORMATION OF ALPHA-BENZYLATED ARIFLO™ (41)

Treatment of compound 40 (prepared according to the literature procedure: *Journal of Medicinal Chemistry*, 1998, 41, 821-835) with LDA in THF at -78°C, followed by 4-(benzyloxy)-3-methoxybenzyl bromide 44 may provide the desired compound 41. In Scheme 10, Ocyclopentyl represents cyclopentyloxy.


Scheme 10

10 Preparation of Compound 41

To a cooled solution of diisopropylamine in anhydrous THF at -78°C was added *n*-butyllithium under argon. The mixture was then stirred at -78°C for 1 hour. To this was added a solution of compound 40 in a mixture of anhydrous THF and 1,3-dimethyl-3,4,5,6-tetrahydro-2(1H)-pyrimidinone. The reaction mixture was stirred at -78°C for 1 hour and then 4-(cyclopentyloxy)-3-methoxybenzyl bromide (Compound 44, prepared according to Scheme 11 below) in anhydrous THF was added in one portion. The resulting mixture was stirred at -78°C for an additional 3 hours, warmed to 0°C, and diluted with 2 N HCl. The aqueous layer was extracted with EtOAc. The combined organic layers were washed with 2 N HCl, followed by brine, and then dried over MgSO₄. The mixture was filtered, and the filtrate was evaporated to dryness. The resulting residue was purified by silica gel column chromatography to afford compound 41.

In Scheme 11, treatment of compound 42 with cyclopentyl bromide, potassium iodide and potassium carbonate in DMF gives the corresponding O-cyclopentyl (OcPent) derivative 43, which is treated with PBr₃ in diethyl ether to give desired bromide compound 44.

111

Scheme 11Synthesis of Compound 43

To a suspension of 4-hydroxy-3-methoxybenzyl alcohol 42 (1.00 g, 6.49 mmol), potassium carbonate (1.79 g, 12.98 mmol) and potassium iodide (29.1 mg, 0.175 mmol) in DMF (10 mL) was slowly added cyclopentyl bromide (0.91 mL, 8.44 mmol). The reaction mixture was stirred at 65° C for 24 hours. After cooling to room temperature, the mixture was diluted with diethyl ether (50 mL) and washed with water (2 x 25 mL). Drying over anhydrous $MgSO_4$, filtration and evaporation of the filtrate *in vacuo* gave crude yellow solid which was purified by silica gel column chromatography (hexanes/EtOAc, 3:1) to give compound 43 (0.502 g, 35%) as a pale yellow solid.

Synthesis of Compound 44


To a solution of compound 43 (0.48 g, 2.17 mmol) in anhydrous diethyl ether (8 mL) was slowly added PBr_3 (0.10 mL, 1.09 mmol) *via* syringe, and the resulting mixture was stirred at room temperature for 2 hours. The mixture was diluted with diethyl ether (50 mL) and washed with saturated aqueous $NaHCO_3$ (2 x 25 mL) and brine (2 x 25 mL). The organic layer was dried over anhydrous $MgSO_4$, and the solvent was removed under reduced pressure to afford compound 44 (0.577 g, 93%) as a white solid.

EXAMPLE 9

FORMATION OF N-BENZYLATED WAY-PDA-641

Exemplary synthetic methodology to introduce a substituted benzyl group into compound 45 is described in Scheme 12. In Scheme 12, compound 45 (*i.e.*, WAY-PDA-641) is treated with NaH in DMF, followed by 3,4-difluorobenzyl bromide to give the desired compound 46 (*i.e.*, benzylated WAY-PDA-641).


112

Scheme 12

EXAMPLE 10

FORMATION OF N-BENZYLATED RO-20-1724

- 5 Exemplary synthetic methodology to introduce a substituted benzyl group into compound 47 is described in Scheme 13. In Scheme 13, compound 47 (*i.e.*, Ro-20-1724) is treated with NaH in DMF, followed by 3,4-difluorobenzyl bromide to give the desired compounds 48 and 49 (*i.e.*, benzylated Ro-20-1724 compounds). Excess NaH and 3,4-difluorobenzyl bromide may be used to provide the compound
- 10 with benzyl groups on both nitrogens.

Scheme 13


EXAMPLE 11

FORMATION OF N-BENZYLATED RP 73401

- 15 Exemplary synthetic methodology to introduce a substituted benzyl group into compound 50 is described in Scheme 14. In Scheme 14, compound 50 (*i.e.*,

RP 73401) is treated with NaH in DMF, followed by 3,4-difluorobenzyl bromide to give the desired compound 51 (*i.e.*, benzylated RP 73401).

Scheme 14



5


EXAMPLE 12

FORMATION OF N-BENZYLATED CP-80,633-A

Exemplary synthetic methodology to introduce a substituted benzyl group into compound 52 is described in Scheme 15. In Scheme 15, compound 52 (*i.e.*, CP-80,633-A) is treated with NaH in DMF, followed by 3,4-difluorobenzyl bromide to give the desired compound 53 (*i.e.*, benzylated CP-80,633-A). Excess NaH and 3,4-difluorobenzyl bromide may be used to provide the compound with benzyl groups on both nitrogens.

10

Scheme 15




114

EXAMPLE 13

FORMATION OF N-BENZYLATED AROFYLLINE™

Exemplary synthetic methodology to introduce a substituted benzyl group into compound **54** is described in Scheme 16. In Scheme 16, compound **54** (AROFYLLINE™) is treated with NaH in DMF, followed by 3,4-difluorobenzyl bromide to give the desired compound **55** (i.e., benzylated AROFYLLINE™).

Scheme 16




EXAMPLE 14

FORMATION OF N-BENZYLATED CIPAMFYLLINE™

Exemplary synthetic methodology to introduce a substituted benzyl group into compound **56** is described in Scheme 17. In Scheme 17, compound **56** (i.e., CIPAMFYLLINE™) is treated with KOtBu in DMF, followed by 3,4-difluorobenzyl bromide to give the desired compound **57** (i.e., benzylated CIPAMFYLLINE™).

Scheme 17



EXAMPLE 15

IN VITRO TESTING FOR EMETOGENIC RESPONSE

Table 1 shows the PDE4 and HARBS activities of compounds of the present invention compared to that of ROLIPRAM™. Compounds 25 and 26 (see Scheme 6) comprise two different derivatives of ROLIPRAM™ in which the substitution pattern of the added benzyl group at the alpha position differs. As these data clearly show, benzylation at the alpha carbon position of ROLIPRAM™ dramatically reduces the HARBS binding affinity of these derivatives by 92 and 56 fold for compound 26 and 25, respectively. In reducing the HARBS binding affinities the PDE4:HARBS ratios for these compounds is also dramatically reduced to 1.1 and 2.5 for compounds 25 and 26 respectively compared to 65 for ROLIPRAM™. This effect on the HARBS affinities and the ratio of PDE4:HARBS is expected to result in reduced emetogenic potential as well as improved therapeutic window for these compounds.

Table 1

EFFECT OF BENZYLATION AT THE ALPHA POSITION OF ROLIPRAM™ ON POTENCY OF PDE4 INHIBITION AND BINDING TO THE HARBS SITE OF PDE4.

Compound	α -substitution (+/- benzyl)	PDE4 IC ₅₀ (μ M)	HARBS IC ₅₀ (μ M)	Ratio (PDE4/HARBS)
ROLIPRAM™	No	0.42	.0065	65.1
26	Yes	1.49	.6012	2.5
25	Yes	0.40	.3608	1.1

These data also show that the nature of the alpha-positioned benzyl substituent influences the degree of reduction in HARBS binding as well as the reduction in potency of PDE4 inhibitory activity. The example of compound 25 also shows that the HARBS binding affinity of ROLIPRAM™ may be reduced without affecting the PDE4 inhibitory activity (and anti-inflammatory efficacy) by the correct choice of benzyl substituent.

Table 2 shows the PDE4 and HARBS activities of a compound of the present invention compared to that of ARIFLO™. Compound 41 is a derivative of ARIFLO™ in which a substituted benzyl group has been added to the carbon alpha to the carboxyl group. The data show that benzylation at the alpha position of ARIFLO™ reduces the HARBS binding affinity of this derivative by 7.9 fold relative to that of the parent compound. In this example the potency in inhibition of PDE4 catalytic activity is

also reduced 7.5 fold for the benzylated derivative resulting in an improvement in the PDE4/HARBS ratio from 2.4 to 2.2.

Table 2

EFFECT OF BENZYLATION AT THE ALPHA POSITION OF ARIFLO™ ON POTENCY OF PDE4
INHIBITION AND BINDING TO THE HARBS SITE OF PDE4.

Compound	α -substitution (+/- benzyl)	PDE4 IC ₅₀ (μ M)	HARBS IC ₅₀ (μ M)	Ratio (PDE4/HARBS)
ARIFLO™	No	0.18	.076	2.4
41	Yes	1.35	.6012	2.2

As demonstrated by the above examples for benzylation of ROLIPRAM™, the nature of the added benzyl group (substitution pattern) may strongly influence the degree of improvement of the PDE4:HARBS ratio and therefore a different benzyl group added to the alpha carbon of ARIFLO™ could confer improved ratios relative to the example above.

Additionally, the site of addition of the benzyl group to the PDE4 inhibitor may strongly influence the degree of improvement (reduction) in the PDE 4:HARBS ratio relative to the non-benzylated compound. As such, addition of a benzyl moiety to the beta or gamma carbons of ARIFLO™ may be desirable to that of alpha substitution as in the above example.

EXAMPLE 16

IN VIVO TESTING FOR EMETOGENIC RESPONSE

The most recognized pre-clinical animal model for emesis is the ferret.

The ferret undergoes a very characteristic, reproducible and well-documented series of behavioral responses, including backward walking, pawing of the mouth, arching of the back, retching and vomiting when dosed with an emetogenic compound. With respect to PDE4 inhibitors their emetogenic potential has been correlated with the ability of compounds to bind to a conformer of the enzyme to which ROLIPRAM™ binds with nanomolar affinity (the HARBS binding site). The central nervous system (CNS) and emetic effects of PDE4 inhibitors are therefore thought in large part to be due to the interaction of these compounds with this particular conformer of PDE4 which is especially prevalent in both the central and peripheral nervous system as well as in cells of the lumen of the gastrointestinal tract (*i.e.*, parietal cells of the stomach).

Benzylation of compounds with PDE4 inhibitory activity (PDE4 inhibitors) according to the present invention has been shown to reduce their HARBS binding potency (see EXAMPLE 15) and thus by logical and rationale extension, based on the existing body of knowledge in the field, to reduce their emetogenic potential. To confirm the in-vitro data of the present invention indicating that incorporation of a benzyl group at the alpha position of ROLIPRAM™ reduces binding to the HARBS conformer of PDE4 compared to that of the parent compound ROLIPRAM™, these two compounds were evaluated for emetogenic potency after intravenous infusion in the ferret model of emesis.

10 Protocol

Adult male ferrets (*Mustela putorius furo*, sable, .75-1.5 kg, Marshall Farms, USA) were quarantined and acclimated to the testing facility for seven days prior to initiation of the study. For intravenous infusion dosing, a catheter was surgically inserted in the jugular vein a minimum of 14 hours prior to administration of test articles and the catheter was maintained for patency by flushing with heparin solution daily. After surgery for catheter insertion the ferrets were individually housed in cages. Animals were fasted 12-16 hours prior to dosing and during the behavioral assessment period. They were returned to normal food diet after the observation periods. Animals found acceptable for inclusion in the study were randomly assigned to the each of the study groups (n=3 for each of the two treatment groups and n=3 for the vehicle negative control group).

Test compounds (ROLIPRAM™ and Compound 25) were freshly dissolved on the day of dosing in the vehicle, polyethylene glycol (PEG) 200 (Sigma catalogue number P-3015), to the required concentration in order to achieve the target dose. Animals in the control (PEG 200) and treatment (ROLIPRAM™ or Compound 25) groups were administered the dosing formulation at 3 ml/kg by intravenous infusion over a 5 minute period via the catheter inserted in the jugular. Continual observation for signs of emetic behavior were begun for each of the animals in the study groups immediately following completion of the 5 minute dosing period and for a period of three hours. Sequential recordings of emetic behaviour were noted on a separate data collection sheet for each animal which included the time of onset of each event as well as the duration and nature of the event (*i.e.*, retching, gagging, vomiting, back arching, backward walking etc.).

The target doses for each of the test articles are shown below in Table 3. If animals expressed no emetic behaviour at the lowest dosage then animals were dosed

again with semi-log increasing doses of test article and the three hour observation period repeated until the emetic threshold (lowest of test doses causing emesis) was reached or until the highest of the 4 target doses was completed. There was a 24 hr. washout period between each of the successive ascending doses of test article to allow the compound to clear the animals' system. For a dose level to be considered at the threshold of emesis at least one of the ferrets must have exhibited definite emetic signs (*i.e.*, retching and/or vomiting). If the emetic response was minimal at a particular dose level then the next dose level could be performed to confirm the threshold level.

Table 3

10 STUDY GROUP DESIGNATIONS FOR DETERMINATION OF EMETIC THRESHOLD IN FERRETS OF ROLIPRAM™ AND BENZYLATED ROLIPRAM™

Treatment	No. of animals	Route	Dosage (mg/kg)
Vehicle (PEG 200)	3	i.v.	-
ROLIPRAM™	3	i.v.	0.03, 0.1, 0.3, 1.0
Compound 25	3	i.v.	0.1, 0.3, 1.0, 3.0

Results

Table 4 shows the dosage required for each of the test compounds (rolipram and compound 25) to achieve a threshold emetic response after 5 minutes intravenous infusion in the ferret. A threshold emetic response was achieved after dosing with 0.3 mg/kg rolipram. At this dosage level of rolipram, one of the three ferrets exhibited a definite emetic response as shown by 5 mild retches observed over a time period of 9 minutes starting at 10 minutes post-dosing. Emetic signs were also evident for rolipram at the previous 0.1 mg/kg dose level.

In contrast to rolipram there was no emetic behaviour whatsoever demonstrated by the ferrets administered compound 25 up to and including the highest dose level tested (3 mg/kg). Thus for compound 25 the emetic threshold in ferrets is greater than 3 mg/kg after intravenous infusion dosing. These data therefore show that at least 10-fold higher doses of the benzylated compound are required to achieve emesis compared to the non-benzylated compound, rolipram. Since these compounds have equivalent PDE4 catalytic inhibitory activity, the benzylated compound in addition to being significantly less emetic than its non-benzylated parent, would be expected to have similar anti-inflammatory efficacy and thus a markedly improved therapeutic ratio.

These in-vivo data demonstrated in the ferret model of emesis, therefore confirm and support the in-vitro biochemical findings presented above that benzylation reduces binding of compounds to the HARBS conformer of PDE4 and thus reduces their emetogenic potential.

5

Table 4

COMPARISON OF THE EMETIC THRESHOLD OF ROLIPRAM AND BENZYLATED ROLIPRAM
AFTER INTRAVENOUS INFUSION IN FERRETS

Treatment	Emetic Threshold (mg/kg)
Rolipram	0.3
Compound 25	>3.0

10 All publications and patent applications mentioned in this specification
are herein incorporated by reference to the same extent as if each individual publication
or patent application was specifically and individually incorporated by reference. For
example, the book in *Comprehensive Organic Transformations, A Guide to Functional
Group Preparations*, Second Edition, Richard C. Larock, John Wiley and Sons, Inc.,
1999, and particularly the references cited therein, is incorporated herein by reference
15 for all purposes.

From the foregoing it will be appreciated that, although specific
embodiments of the invention have been described herein for purposes of illustration,
various modifications may be made without deviating from the spirit and scope of the
invention. Accordingly, the invention is not limited except as by the appended claims.

CLAIMS

1. A method for improving the therapeutic ratio of PDE4 inhibitors through reduction or elimination of emetogenic properties while allowing retention of PDE activity, the method comprising benzylation of a PDE4 inhibitor, wherein the benzylation places a benzyl group on a carbon atom of a PDE4 inhibitor, to provide a benzylated PDE4 inhibitor.


2. The method according to claim 1 wherein the PDE4 inhibitor is selected from ROLIPRAM™ and ARIFLO™.

3. The method of claim 1 wherein the benzylation replaces an extractable hydrogen of the PDE4 inhibitor with a benzyl group.

4. The method of claim 1 wherein the PDE4 inhibitor comprises a carbonyl group, and the benzylation replaces an extractable hydrogen of the PDE4 that is at a location alpha to the carbonyl group.

5. The method of claim 1 wherein the PDE4 inhibitor is reacted with a base followed by reaction with a benzylating agent.

6. The method of claim 1 wherein the benzylated PDE4 inhibitor comprises at least one benzyl group (Bzl) of the formula




wherein each of the numerals 1, 2, 3, 4 and 5 may be carbon or nitrogen, with the proviso that the ring is aromatic and contains at least 3 carbon atoms;

n is 5 and R^1 at each occurrence is independently selected from halogen, nitro, R^2 , $NR^2_{(m)}$, and OR^2 wherein $m = 0, 1, 2$, or 3 and R^2 at each occurrence is independently selected from H, C_1 - C_8 alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C_3 - C_7 cycloalkyl, C_6 - C_9 polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxyl, alkylcarboxylate, carboxylate, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion,

phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R² group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-C₈alkoxy; and the aryl portion of an R² group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen.


7. The method of claim 1 wherein the benzylated PDE4 inhibitor comprises at least one benzyl group (Bzl) of the formula



wherein each of the numerals 1, 2, 3, 4 and 5 may be carbon or nitrogen, with the proviso that the ring is aromatic and contains at least 4 carbon atoms;

n is 5 and R¹ at each occurrence is independently selected from halogen, nitro, R², and OR² wherein R² at each occurrence is independently selected from H, C₁-C₈alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C₃-C₇cycloalkyl, C₆-C₉polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, phenyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R² group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-C₈alkoxy; and the phenyl portion of an R² group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen.

8. The method of claim 1 wherein the benzylated PDE4 inhibitor comprises at least one benzyl group (Bzl) of the formula




wherein each of the numerals 1, 2, 3, 4 and 5 may be carbon or nitrogen, with the proviso that the ring is aromatic and contains at least 3 carbon atoms;

n is 5 and R¹ at each occurrence is independently selected from halogen, nitro, R², NR²_(m), and OR² wherein m = 0,1,2, or 3 and R² at each occurrence is independently

selected from H, C₁-C₈alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C₃-C₇cycloalkyl, C₆-C₉polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R² group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-C₈alkoxy; and the aryl portion of an R² group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen;

with the proviso that either (a) at least one R¹ is OR², or (b) the benzyl ring contains at least two R¹ groups that are not hydrogen.

9. The method of claim 1 wherein the benzylated PDE4 inhibitor comprises at least one benzyl group (Bzl) of the formula




wherein each of the numerals 1, 2, 3, 4 and 5 may be carbon or nitrogen, with the proviso that the ring is aromatic and contains at least 3 carbon atoms;

n is 5 and R¹ at each occurrence is independently selected from halogen, nitro, R², NR²_(m), and OR² wherein m = 0,1,2, or 3 and R² at each occurrence is independently selected from H, C₁-C₈alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C₃-C₇cycloalkyl, C₆-C₉polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R² group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-C₈alkoxy; and the aryl portion of an R² group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen;

with the proviso that R₁ is hydrogen at either 1, 2, or 3 occurrences of n, but not 4 or 5 occurrences.

10. The method of claim 1 wherein the benzylated PDE4 inhibitor comprises a benzyl group, and independently of the benzyl group the benzylated PDE4 inhibitor further comprises the group




wherein p is 5; and R^3 at each occurrence is independently selected from halogen, nitro, R^4 , $NR^4_{(q)}$, and OR^4 wherein $q = 0, 1, 2$, or 3 and R^4 at each occurrence is independently selected from H, C_1 - C_8 alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C_3 - C_7 cycloalkyl, C_6 - C_9 polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C_1 - C_4 alkyl and indanyl; wherein the alkyl portion of an R^4 group may be optionally substituted with one or more fluorine atoms, hydroxyl or C_1 - C_8 alkoxy; and the aryl portion of an R^4 group may be optionally substituted with C_1 - C_8 alkyl, C_1 - C_8 alkoxy or halogen;

with the proviso that R^3 at the para position relative to A is alkoxy, and R^3 at one meta position is selected from alkyl having at least 3 carbons and alkoxy having at least 3 carbons.

11. The method of claim 10 wherein R^3 is OR^4 in at least one occurrence wherein R^4 is selected from C_1 - C_8 alkyl, C_3 - C_7 cycloalkyl, and C_6 - C_9 polycycloalkyl.

12. The method of claim 1 wherein the PDE4 inhibitor is ROLIPRAM™ or an analog thereof.

13. The method of claim 12 wherein the benzylated ROLIPRAM™ or analog thereof has the formula:



and isolated stereoisomers, salts and solvates thereof, wherein

Bzl is a benzyl group;


A is selected from a direct bond, optionally substituted C₁-C₅alkylene, optionally substituted C₂-C₅alkenyl and optionally substituted phenylene;

p is 5; and R³ at each occurrence is independently selected from halogen, nitro, R⁴, NR⁴_(q), and OR⁴ wherein q = 0,1,2, or 3 and R⁴ at each occurrence is independently selected from H, C₁-C₈alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C₃-C₇cycloalkyl, C₆-C₉polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R⁴ group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-C₈alkoxy; and the aryl portion of an R⁴ group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen.

125


14. The method of claim 13 wherein A is a direct bond; and in at least one occurrence, R^3 is OR^4 wherein R^4 is selected from C_1 - C_8 alkyl, C_3 - C_7 cycloalkyl, and C_6 - C_9 polycycloalkyl.

15. The method of claim 14 wherein the benzylated ROLIPRAM™ or analog thereof has the formula:



16. The method of claim 1 wherein the PDE4 inhibitor is ARIFLO™ or an analog thereof.

17. The method of claim 16 wherein the benzylated ARIFLO™ or analog thereof has the formula



and isolated stereoisomers, salts and solvates thereof, wherein

Bzl is a benzyl group;

p is 5; and R^3 at each occurrence is independently selected from halogen, nitro, R^4 , $NR^4_{(q)}$, and OR^4 wherein $q = 0, 1, 2$, or 3 and R^4 at each occurrence is independently selected from H , C_1 - C_8 alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C_3 - C_7 cycloalkyl, C_6 - C_9 polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl,

126


heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R⁴ group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-C₈alkoxy; and the aryl portion of an R⁴ group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen;

R⁵ is selected from H, halogen, cyano, C₁-C₈alkyl, and C₁-C₈alkoxy; and


R⁶ is selected from H, positively charged species, and C₁-C₈alkyl.

18. The method of claim 17 wherein
in at least one occurrence, R³ is OR⁴ wherein R⁴ is selected from C₁-C₈alkyl, C₃-C₇cycloalkyl, and C₅-C₉bicycloalkyl; and
R⁶ is selected from H and positively charged species.

19. The method of claim 18 wherein the benzylated ARIFLO™ or analog thereof has the formula



20. The method of claim 16 wherein the benzylated ARIFLO™ or analog thereof has the formula



and isolated stereoisomers, salts and solvates thereof, wherein

Bzl is a benzyl group;

p is 5; and R³ at each occurrence is independently selected from halogen, nitro, R⁴, NR⁴_(q), and OR⁴ wherein q = 0,1,2, or 3 and R⁴ at each occurrence is independently

selected from H, C₁-C₈alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C₃-C₇cycloalkyl, C₆-C₉polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R⁴ group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-C₈alkoxy; and the aryl portion of an R⁴ group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen;

R⁵ is selected from H, halogen, cyano, C₁-C₈alkyl, and C₁-C₈alkoxy; and


R⁶ is selected from H, positively charged species, and C₁-C₈alkyl.

21. The method of claim 20 wherein


in at least one occurrence, R³ is OR⁴ wherein R⁴ is selected from C₁-C₈alkyl, C₃-C₇cycloalkyl, and C₅-C₉bicycloalkyl; and

R⁶ is selected from H and positively charged species.

22. The method of claim 21 wherein the benzylated ARIFLO™ or analog thereof has the formula



23. A method for improving the therapeutic ratio of PDE4 inhibitors through reduction or elimination of emetogenic properties while allowing retention of PDE activity, the method comprising benzylation of a PDE4 inhibitor, wherein the benzylation places a benzyl group (Bzl) on a nitrogen atom of a PDE4 inhibitor, to provide a benzylated PDE4 inhibitor, wherein the benzylated PDE4 inhibitor comprises at least one benzyl group (Bzl) of the formula



wherein each of the numerals 1, 2, 3, 4 and 5 may be carbon or nitrogen, with the proviso that the ring is aromatic and contains at least 3 carbon atoms;

n is 5 and R^1 at each occurrence is independently selected from halogen and OR^2 wherein R^2 at each occurrence is independently selected from H and C_1-C_8 alkyl; with the provisos that the benzyl ring contain no more than three hydrogen substituents and no more than one methoxy substituent.

24. The method of claim 23 wherein each of the numerals 1, 2, 3, 4 and 5 is carbon.


25. The method of claim 23 wherein Bzl has two halogen substituents on the aromatic ring.

26. The method according to claim 23 wherein the PDE4 inhibitor is selected from ROLIPRAM™, WAY-PDA-641, Ro-20-1724, RP 73401, CP-80,633-A, AROFYLLINE™ and CIPAMFYLLINE™.

27. The method of claim 23 wherein the benzylation replaces an extractable hydrogen of the PDE4 inhibitor a benzyl group.

28. The method of claim 23 wherein the PDE4 inhibitor is reacted with a base followed by a benzylating agent.

29. The method of claim 23 wherein the benzylated ROLIPRAM™ or analog thereof has the formula:




and isolated stereoisomers, salts and solvates thereof, wherein Bzl is a benzyl group;

A is selected from a direct bond, optionally substituted C₁-C₃alkylene, optionally substituted C₂-C₅alkenyl and optionally substituted phenylene;

p is 5 and R³ at each occurrence is independently selected from halogen, nitro, R⁴, NR⁴_(q), and OR⁴ wherein q = 0,1,2, or 3 and R⁴ at each occurrence is independently selected from H, C₁-C₈alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C₃-C₇cycloalkyl, C₆-C₉polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R⁴ group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-C₈alkoxy; and the aryl portion of an R⁴ group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen.


30. The method of claim 29 wherein A is a direct bond; p is 1, 2, or 3; and in at least one occurrence, R³ is OR⁴ wherein R⁴ is selected from C₁-C₈alkyl, C₃-C₇cycloalkyl, and C₆-C₉polycycloalkyl.

31. The method of claim 30 wherein the benzylated ROLIPRAM™ or analog thereof has the formula



32. The method of claim 23 wherein the PDE4 inhibitor is WAY-PDA-641 or an analog thereof.

33. The method of claim 32 wherein the benzylated WAY-PDA-641 or analog thereof has the formula:



and isolated stereoisomers, salts and solvates thereof, wherein


Bzl is a benzyl group;

p is 5 and R^3 at each occurrence is independently selected from halogen, nitro, R^4 , $NR^4_{(q)}$, and OR^4 wherein $q = 0, 1, 2$, or 3 and R^4 at each occurrence is independently selected from H, C_1 - C_8 alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C_3 - C_7 cycloalkyl, C_6 - C_9 polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C_1 - C_4 alkyl and indanyl; wherein the alkyl portion of an R^4 group may be optionally substituted with one or more fluorine atoms, hydroxyl or C_1 - C_8 alkoxy; and the aryl portion of an R^4 group may be optionally substituted with C_1 - C_8 alkyl, C_1 - C_8 alkoxy or halogen.

34. The method of claim 33 wherein p is 1, 2, or 3; and in at least one occurrence, R^3 is OR^4 wherein R^4 is selected from C_1 - C_8 alkyl, C_3 - C_7 cycloalkyl, and C_6 - C_9 polycycloalkyl.


131

35. The method of claim 33 wherein the benzylated WAY-PDA-641 or analog thereof has the formula:



36. The method of claim 23 wherein the PDE4 inhibitor is Ro-20-1724 or an analog thereof.

37. The method of claim 36 wherein the benzylated Ro-20-1724 or analog thereof has the formula:



and isolated stereoisomers, salts and solvates thereof, wherein

Bzl is a benzyl group;


p is 5 and R^3 at each occurrence is independently selected from halogen, nitro, R^4 , $NR^4_{(q)}$, and OR^4 wherein $q = 0, 1, 2$, or 3 and R^4 at each occurrence is independently selected from H, C_1 - C_8 alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C_3 - C_7 cycloalkyl, C_6 - C_9 polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C_1 - C_4 alkyl and indanyl; wherein the alkyl portion of an R^4 group may be optionally substituted

132


with one or more fluorine atoms, hydroxyl or C₁-C₈alkoxy; and the aryl portion of an R⁴ group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen.

38. The method of claim 34 wherein p is 1, 2, or 3; and in at least one occurrence, R³ is OR⁴ wherein R⁴ is selected from C₁-C₈alkyl, C₃-C₇cycloalkyl, and C₆-C₉polycycloalkyl.

39. The method of claim 37 wherein the benzylated Ro-20-1724 or analog thereof has the formula:



40. The method of claim 36 wherein the benzylated Ro-20-1724 or analog thereof has the formula:



and isolated stereoisomers, salts and solvates thereof, wherein


Bzl is a benzyl group;

p is 5 and R³ at each occurrence is independently selected from halogen, nitro, R⁴, NR⁴_(q), and OR⁴ wherein q = 0,1,2, or 3 and R⁴ at each occurrence is independently selected from H, C₁-C₈alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C₃-C₇cycloalkyl, C₆-C₉polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate,

alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R⁴ group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-C₈alkoxy; and the aryl portion of an R⁴ group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen.


41. The method of claim 40 wherein p is 1, 2, or 3; and in at least one occurrence, R³ is OR⁴ wherein R⁴ is selected from C₁-C₈alkyl, C₃-C₇cycloalkyl, and C₆-C₉polycycloalkyl.

42. The method of claim 41 wherein the benzylated Ro-20-1724 or analog thereof has the formula:



43. The method of claim 23 wherein the PDE4 inhibitor is RP 73401 or an analog thereof.

44. The method of claim 43 wherein the benzylated RP 73401 or analog thereof has the formula:



and isolated stereoisomers, salts and solvates thereof, wherein


Bzl is a benzyl group;

p is 5 and R^3 at each occurrence is independently selected from halogen, nitro, R^4 , $NR^4_{(q)}$, and OR^4 wherein $q = 0, 1, 2$, or 3 and R^4 at each occurrence is independently selected from H, C_1 - C_8 alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C_3 - C_7 cycloalkyl, C_6 - C_9 polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C_1 - C_4 alkyl and indanyl; wherein the alkyl portion of an R^4 group may be optionally substituted with one or more fluorine atoms, hydroxyl or C_1 - C_8 alkoxy; and the aryl portion of an R^4 group may be optionally substituted with C_1 - C_8 alkyl, C_1 - C_8 alkoxy or halogen.

45. The method of claim 44 wherein p is 1, 2, or 3; and in at least one occurrence, R^3 is OR^4 wherein R^4 is selected from C_1 - C_8 alkyl, C_3 - C_7 cycloalkyl, and C_6 - C_9 polycycloalkyl.


135

46. The method of claim 44 wherein the benzylated RP 73401 or analog thereof has the formula:



47. The method of claim 23 wherein the PDE4 inhibitor is CP-80,633-A or an analog thereof.

48. The method of claim 47 wherein the benzylated CP-80,633-A or analog thereof has the formula:



and isolated stereoisomers, salts and solvates thereof, wherein

Bzl is a benzyl group;


p is 5 and R^3 at each occurrence is independently selected from nitro, R^4 , $NR^4_{(q)}$, and OR^4 wherein $q = 0, 1, 2$, or 3 and R^4 at each occurrence is independently selected from H, C_1 - C_8 alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C_3 - C_7 cycloalkyl, C_6 - C_9 polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in

136

the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R⁴ group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-C₈alkoxy; and the aryl portion of an R⁴ group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen.


49. The method of claim 48 wherein p is 1, 2, or 3; and in at least one occurrence, R³ is OR⁴ wherein R⁴ is selected from C₁-C₈alkyl, C₃-C₇cycloalkyl, and C₆-C₉polycycloalkyl.

50. The method of claim 49 wherein the benzylated CP-80,633-A or analog thereof has the formula:



51. The method of claim 23 wherein the PDE4 inhibitor is AROFYLLINE™ or an analog thereof.

52. The method of claim 51 wherein the benzylated AROFYLLINE™ or analog thereof has the formulae:




and isolated stereoisomers, salts and solvates thereof, wherein

Bzl is a benzyl group;

p is 5 and R^3 at each occurrence is independently selected from halogen, nitro, R^4 , $NR^4_{(q)}$, and OR^4 wherein $q = 0, 1, 2$, or 3 and R^4 at each occurrence is independently selected from H, C_1 - C_8 alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C_3 - C_7 cycloalkyl, C_6 - C_9 polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C_1 - C_4 alkyl and indanyl; wherein the alkyl portion of an R^4 group may be optionally substituted with one or more fluorine atoms, hydroxyl or C_1 - C_8 alkoxy; and the aryl portion of an R^4 group may be optionally substituted with C_1 - C_8 alkyl, C_1 - C_8 alkoxy or halogen.


53. The method of claim 52 wherein the benzylated AROFYLLINE™ or analog thereof has the formula




138

55. The method of claim 52 wherein p is 1, 2 or 3; and in at least one occurrence, R^3 is halogen or OR^4 wherein R^4 is selected from C_1 - C_8 alkyl, C_3 - C_7 cycloalkyl, and C_6 - C_9 polycycloalkyl.

56. The method of claim 52 wherein the benzylated AROFYLLINE™ or analog thereof has the formula:




57. The method of claim 52 wherein the benzylated AROFYLLINE™ or analog thereof has the formula:



58. The method of claim 23 wherein the PDE4 inhibitor is CIPAMFYLLINE™ or an analog thereof.


130

59. The method of claim 58 wherein the benzylated CIPAMFYLLINE™ or analog thereof has the formulae:




and isolated stereoisomers, salts and solvates thereof, wherein Bzl is a benzyl group.

60. A C-benzylated ROLIPRAM™ or analog thereof compound of the formula



and isolated stereoisomers, salts and solvates thereof, wherein Bzl is a benzyl group of the formula



wherein each of the numerals 1, 2, 3, 4 and 5 may be carbon or nitrogen, with the proviso that the ring is aromatic and contains at least 3 carbon atoms;

n is 5 and R¹ at each occurrence is independently selected from halogen, nitro, R², NR²_(m), and OR² wherein m = 0, 1, 2, or 3 and R² at each occurrence is independently selected from H, C₁-C₈alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C₃-C₇cycloalkyl, C₆-C₉polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate,

alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R² group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-C₈alkoxy; and the aryl portion of an R² group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen;


A is selected from a direct bond, optionally substituted C₁-C₅alkylene, optionally substituted C₂-C₅alkenyl and optionally substituted phenylene;

p is 5 and R³ at each occurrence is independently selected from halogen, nitro, R⁴, NR⁴_(q), and OR⁴ wherein q = 0,1,2, or 3 and R⁴ at each occurrence is independently selected from H, C₁-C₈alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C₃-C₇cycloalkyl, C₆-C₉polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R⁴ group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-C₈alkoxy; and the aryl portion of an R⁴ group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen;


with the proviso that R³ at the para position relative to A is alkoxy, and R³ at one meta position is selected from alkyl having at least 3 carbons and alkoxy having at least 3 carbons.

61. The compound of claim 60 wherein A is a direct bond; and in at least one occurrence, R³ is OR⁴ wherein R² is selected from C₁-C₈alkyl, C₃-C₇cycloalkyl, and C₆-C₉polycycloalkyl.


62. The compound of claim 60 having the formula



63. An N-benzylated ROLIPRAM™ or analog thereof compound of the formulae



and isolated stereoisomers, salts and solvates thereof, wherein Bzl is a benzyl group of the formula



wherein each of the numerals 1, 2, 3, 4 and 5 may be carbon or nitrogen, with the proviso that the ring is aromatic and contains at least 3 carbon atoms;


n is 5 and R^1 at each occurrence is independently selected from halogen, nitro, R^2 , $NR^2_{(m)}$, and OR^2 wherein $m = 0, 1, 2$, or 3 and R^2 at each occurrence is independently selected from H, C_1 - C_8 alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C_3 - C_7 cycloalkyl, C_6 - C_9 polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C_1 - C_4 alkyl and indanyl; wherein the alkyl portion of an R^2 group may be optionally substituted with one or more fluorine atoms, hydroxyl or C_1 - C_8 alkoxy; and the aryl portion of an R^2 group may be optionally substituted with C_1 - C_8 alkyl, C_1 - C_8 alkoxy or halogen;

A is selected from a direct bond, optionally substituted C_1 - C_5 alkylene, optionally substituted C^2 - C^5 alkenyl and optionally substituted phenylene;


p is 5 and R^3 at each occurrence is independently selected from halogen, nitro, R^4 , $NR^4_{(q)}$, and OR^4 wherein $q = 0, 1, 2$, or 3 and R^4 at each occurrence is independently selected from H, C_1 - C_8 alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C_3 - C_7 cycloalkyl, C_6 - C_9 polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate,

alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R⁴ group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-C₈alkoxy; and the aryl portion of an R⁴ group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen; with the provisos that, at the meta position relative to the A group on the R³-substituted phenyl ring, R³ is alkoxy having at least three carbon atoms, excluding cyclopentyloxy, and at the para position R¹ is not hydrogen.

64. An N-benzylated ROLIPRAM™ or analog thereof compound of the formulae



and isolated stereoisomers, salts and solvates thereof, wherein Bzl is a benzyl group of the formula



wherein each of the numerals 1, 2, 3, 4 and 5 may be carbon or nitrogen, with the proviso that the ring is aromatic and contains at least 3 carbon atoms;

n is 5 and R¹ at each occurrence is independently selected from halogen and OR² wherein R² at each occurrence is independently selected from H and C₁-C₈alkyl; with the provisos that the benzyl ring contain no more than three hydrogen substituents and no more than one methoxy substituent;


A is selected from a direct bond, optionally substituted C₁-C₅alkylene, optionally substituted C²-C⁵alkenyl and optionally substituted phenylene;

p is 5 and R³ at each occurrence is independently selected from halogen, nitro, R⁴, NR⁴_(q), and OR⁴ wherein q = 0,1,2, or 3 and R⁴ at each occurrence is independently


selected from H, C₁-C₈alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C₃-C₇cycloalkyl, C₆-C₉polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R⁴ group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-C₈alkoxy; and the aryl portion of an R⁴ group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen.

65. The compound of claims 63 or 64 wherein A is a direct bond; and in at least one occurrence, R³ is OR⁴ wherein R⁴ is selected from C₁-C₈alkyl, C₃-C₇cycloalkyl, and C₆-C₉polycycloalkyl.


66. The compound of claim 64 having the formula



67. A benzylated ARIFLO™ or analog thereof compound of the formulae



and isolated stereoisomers, salts and solvates thereof, wherein Bzl is a benzyl group of the formula



wherein each of the numerals 1, 2, 3, 4 and 5 may be carbon or nitrogen, with the proviso that the ring is aromatic and contains at least 3 carbon atoms;

n is 5 and R^1 at each occurrence is independently selected from halogen, nitro, R^2 , $NR^2_{(m)}$, and OR^2 wherein $m = 0, 1, 2$, or 3 and R^2 at each occurrence is independently selected from H, C_1 - C_8 alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C_3 - C_7 cycloalkyl, C_6 - C_9 polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C_1 - C_4 alkyl and indanyl; wherein the alkyl portion of an R^2 group may be optionally substituted with one or more fluorine atoms, hydroxyl or C_1 - C_8 alkoxy; and the aryl portion of an R^2 group may be optionally substituted with C_1 - C_8 alkyl, C_1 - C_8 alkoxy or halogen;


p is 5; and R^3 at each occurrence is independently selected from halogen, nitro, R^4 , $NR^4_{(q)}$, and OR^4 wherein $q = 0, 1, 2$, or 3 and R^4 at each occurrence is independently selected from H, C_1 - C_8 alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C_3 - C_7 cycloalkyl, C_6 - C_9 polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C_1 - C_4 alkyl and indanyl; wherein the alkyl portion of an R^4 group may be optionally substituted with one or more fluorine atoms, hydroxyl or C_1 - C_8 alkoxy; and the aryl portion of an R^4 group may be optionally substituted with C_1 - C_8 alkyl, C_1 - C_8 alkoxy or halogen;

R^5 is selected from H, halogen, cyano, C_1 - C_8 alkyl, and C_1 - C_8 alkoxy; and


R^6 is selected from H, positively charged species, and C_1 - C_8 alkyl.

145

68. The compound of claim 67 having the formula




69. The compound of claim 67 having the formula




70. A compound of claim 67 wherein in at least one occurrence, R^3 is OR^4 , wherein R^4 is selected from $\text{C}_1\text{-C}_8$ alkyl, $\text{C}_3\text{-C}_7$ cycloalkyl, and $\text{C}_5\text{-C}_9$ bicycloalkyl; and R^6 is selected from H and positively charged species.

71. The compound of claim 70 having the formula




146


72. The compound of claim 70 having the formula



73. A benzylated WAY-PDA-641 or analog thereof compound of the formula:



and isolated stereoisomers, salts and solvates thereof, wherein Bzl is a benzyl group of the formula




wherein each of the numerals 1, 2, 3, 4 and 5 may be carbon or nitrogen, with the proviso that the ring is aromatic and contains at least 3 carbon atoms;

n is 5 and R^1 at each occurrence is independently selected from halogen, nitro, R^2 , $NR^2_{(m)}$, and OR^2 wherein $m = 0, 1, 2$, or 3 and R^2 at each occurrence is independently selected from H, C_1 - C_8 alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C_3 - C_7 cycloalkyl, C_6 - C_9 polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C_1 - C_4 alkyl and indanyl; wherein the alkyl portion of an R^2 group may be optionally substituted with one or more fluorine atoms, hydroxyl or C_1 - C_8 alkoxy; and the aryl portion of an R^2 group may be optionally substituted with C_1 - C_8 alkyl, C_1 - C_8 alkoxy or halogen;


p is 5; and R^3 at each occurrence is independently selected from halogen, nitro, R^4 , $NR^4_{(q)}$, and OR^4 wherein q = 0,1,2, or 3 and R^4 at each occurrence is independently selected from H, C_1 - C_8 alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C_3 - C_7 cycloalkyl, C_6 - C_9 polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C_1 - C_4 alkyl and indanyl; wherein the alkyl portion of an R^4 group may be optionally substituted with one or more fluorine atoms, hydroxyl or C_1 - C_8 alkoxy; and the aryl portion of an R^4 group may be optionally substituted with C_1 - C_8 alkyl, C_1 - C_8 alkoxy or halogen.

74. The compound of claim 73 wherein in at least one occurrence, R^3 is OR^4 wherein R^4 is selected from C_1 - C_8 alkyl, C_3 - C_7 cycloalkyl, and C_6 - C_9 polycycloalkyl.


75. The compound of claim 74 having the formula:



76. A benzylated Ro-20-1724 or analog thereof compound of the formulae:



and isolated stereoisomers, salts and solvates thereof, wherein Bzl is a benzyl group of the formula



wherein each of the numerals 1, 2, 3, 4 and 5 may be carbon or nitrogen, with the proviso that the ring is aromatic and contains at least 3 carbon atoms;


n is 5 and R^1 at each occurrence is independently selected from halogen, nitro, R^2 , $NR^2_{(m)}$, and OR^2 wherein $m = 0, 1, 2$, or 3 and R^2 at each occurrence is independently selected from H, C_1 - C_8 alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C_3 - C_7 cycloalkyl, C_6 - C_9 polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C_1 - C_4 alkyl and indanyl; wherein the alkyl portion of an R^2 group may be optionally substituted with one or more fluorine atoms, hydroxyl or C_1 - C_8 alkoxy; and the aryl portion of an R^2 group may be optionally substituted with C_1 - C_8 alkyl, C_1 - C_8 alkoxy or halogen;

p is 5; and R^3 at each occurrence is independently selected from halogen, nitro, R^4 , $NR^4_{(q)}$, and OR^4 wherein $q = 0, 1, 2$, or 3 and R^4 at each occurrence is independently selected from H, C_1 - C_8 alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C_3 - C_7 cycloalkyl, C_6 - C_9 polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C_1 - C_4 alkyl and indanyl; wherein the alkyl portion of an R^4 group may be optionally substituted with one or more fluorine atoms, hydroxyl or C_1 - C_8 alkoxy; and the aryl portion of an R^4 group may be optionally substituted with C_1 - C_8 alkyl, C_1 - C_8 alkoxy or halogen;


with the proviso that, relative to the methylene bridge to the heterocyclic ring, R^3 at the para position is alkoxy, and R^3 at a meta position is alkoxy having at least three carbon atoms.

149

77. The compound of claim 76 having the formula




78. The compound of claim 76 having the formula




79. The compound of claim 76 wherein in at least one occurrence, R^3 is OR^4 wherein R^4 is selected from C_1 - C_8 alkyl, C_3 - C_7 cycloalkyl, and C_5 - C_9 bicycloalkyl.

80. The compound of claim 79 having the formula:




150


81. The compound of claim 79 having the formula:



82. A benzylated RP 73401 or analog thereof compound of the formula:



and isolated stereoisomers, salts and solvates thereof, wherein
Bzl is a benzyl group of the formula



wherein each of the numerals 1, 2, 3, 4 and 5 may be carbon or nitrogen, with the proviso that the ring is aromatic and contains at least 3 carbon atoms;


n is 5 and R^1 at each occurrence is independently selected from halogen, nitro, R^2 , $NR^2_{(m)}$, and OR^2 wherein $m = 0, 1, 2$, or 3 and R^2 at each occurrence is independently selected from H, C_1 - C_8 alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C_3 - C_7 cycloalkyl, C_6 - C_9 polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl

having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R² group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-C₈alkoxy; and the aryl portion of an R² group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen;


p is 5; and R³ at each occurrence is independently selected from halogen, nitro, R⁴, NR⁴_(q), and OR⁴ wherein q = 0,1,2, or 3 and R⁴ at each occurrence is independently selected from H, C₁-C₈alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C₃-C₇cycloalkyl, C₆-C₉polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R⁴ group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-C₈alkoxy; and the aryl portion of an R⁴ group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen.

83. The compound of claim 82 wherein in at least one occurrence, R³ is OR⁴ wherein R⁴ is selected from C₁-C₈alkyl, C₃-C₇cycloalkyl, and C₆-C₉polycycloalkyl.


84. The compound of claim 83 wherein the compound has the formula:



85. A benzylated CP-80,633-A or analog thereof compound of the formulae:



and isolated stereoisomers, salts and solvates thereof, wherein Bzl is a benzyl group of the formula



wherein each of the numerals 1, 2, 3, 4 and 5 may be carbon or nitrogen, with the proviso that the ring is aromatic and contains at least 3 carbon atoms;


n is 5 and R^1 at each occurrence is independently selected from halogen, nitro, R^2 , $NR^2_{(m)}$, and OR^2 wherein $m = 0, 1, 2$, or 3 and R^2 at each occurrence is independently selected from H, C_1 - C_8 alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C_3 - C_7 cycloalkyl, C_6 - C_9 polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C_1 - C_4 alkyl and indanyl; wherein the alkyl portion of an R^2 group may be optionally substituted with one or more fluorine atoms, hydroxyl or C_1 - C_8 alkoxy; and the aryl portion of an R^2 group may be optionally substituted with C_1 - C_8 alkyl, C_1 - C_8 alkoxy or halogen;

p is 5; and R^3 at each occurrence is independently selected from nitro, R^4 , $NR^4_{(q)}$, and OR^4 wherein $q = 0, 1, 2$, or 3 and R^4 at each occurrence is independently selected from H, C_1 - C_8 alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C_3 - C_7 cycloalkyl, C_6 - C_9 polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in


the alkyl portion and the amino may be optionally substituted with C_1 - C_4 alkyl and indanyl; wherein the alkyl portion of an R^4 group may be optionally substituted with one or more fluorine atoms, hydroxyl or C_1 - C_8 alkoxy; and the aryl portion of an R^4 group may be optionally substituted with C_1 - C_8 alkyl, C_1 - C_8 alkoxy or halogen; with the proviso that not more than 4 of the R^3 groups are hydrogen.

86. The compound of claim 85 wherein in at least one occurrence, R^3 is OR^4 wherein R^4 is selected from C_1 - C_8 alkyl, C_3 - C_7 cycloalkyl, and C_6 - C_9 polycycloalkyl.


87. The compound of claim 86 wherein the compound has the formula:



88. A benzylated AROFYLLINE™ or analog thereof of the formulae:



and isolated stereoisomers, salts and solvates thereof, wherein Bzl is a benzyl group of the formula




wherein each of the numerals 1, 2, 3, 4 and 5 may be carbon or nitrogen, with the proviso that the ring is aromatic and contains at least 3 carbon atoms;

n is 5 and R^1 at each occurrence is independently selected from halogen, nitro, R^2 , $NR^2_{(m)}$, and OR^2 wherein $m = 0, 1, 2$, or 3 and R^2 at each occurrence is independently selected from H, C_1 - C_8 alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C_3 - C_7 cycloalkyl, C_6 - C_9 polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C_1 - C_4 alkyl and indanyl; wherein the alkyl portion of an R^2 group may be optionally substituted with one or more fluorine atoms, hydroxyl or C_1 - C_8 alkoxy; and the aryl portion of an R^2 group may be optionally substituted with C_1 - C_8 alkyl, C_1 - C_8 alkoxy or halogen;


p is 5; and R^3 at each occurrence is independently selected from halogen, nitro, R^4 , $NR^4_{(q)}$, and OR^4 wherein $q = 0, 1, 2$, or 3 and R^4 at each occurrence is independently selected from H, C_1 - C_8 alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C_3 - C_7 cycloalkyl, C_6 - C_9 polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C_1 - C_4 alkyl and indanyl; wherein the alkyl portion of an R^4 group may be optionally substituted with one or more fluorine atoms, hydroxyl or C_1 - C_8 alkoxy; and the aryl portion of an R^4 group may be optionally substituted with C_1 - C_8 alkyl, C_1 - C_8 alkoxy or halogen.

89. A benzylated AROFYLLINE™ or analog thereof compound of claim 88 having the formula:




155

90. A benzylated AROFYLLINE™ or analog thereof compound of claim 88 having the formula:




91. The compound of claim 87 wherein in at least one occurrence, R^3 is halogen or OR^4 wherein R^4 is selected from C_1 - C_8 alkyl, C_3 - C_7 cycloalkyl, and C_6 - C_9 polycycloalkyl.

92. The compound of claim 91 wherein the compound has the formula:




156


93. The compound of claim 91 wherein the compound has the formula:



94. A benzylated CIPAMFYLLINE™ or analog thereof compound of the formula:



and isolated stereoisomers, salts and solvates thereof, wherein Bzl is a benzyl group of the formula




wherein each of the numerals 1, 2, 3, 4 and 5 may be carbon or nitrogen, with the proviso that the ring is aromatic and contains at least 3 carbon atoms;

n is 5 and R^1 at each occurrence is independently selected from halogen, nitro, R^2 , $NR^2_{(m)}$, and OR^2 wherein $m = 0, 1, 2$, or 3 and R^2 at each occurrence is independently selected from H, C_1 - C_8 alkyl, alkoxyalkyl having 3-7 carbons in the alkoxy portion and 2-4 carbons in the alkyl portion, phenoxyalkyl having 2-6 carbons in the alkyl portion, C_3 - C_7 cycloalkyl, C_6 - C_9 polycycloalkyl, alkylcycloalkyl, hydroxyalkyl, carboxylate, alkylcarboxylate, carboxyl, alkyl carboxyl, amide, alkylamide, aryl, heteroaryl, heteroalkyl, heterocycloalkyl, phenylalkyl having 1-8 carbons in the alkyl portion, phenylaminoalkyl


157

having 2-6 carbons in the alkyl portion and the amino may be optionally substituted with C₁-C₄alkyl and indanyl; wherein the alkyl portion of an R² group may be optionally substituted with one or more fluorine atoms, hydroxyl or C₁-C₈alkoxy; and the aryl portion of an R² group may be optionally substituted with C₁-C₈alkyl, C₁-C₈alkoxy or halogen.

95. A benzylated CIPAMFYLLINE™ or analog thereof compound according to claim 94 of the formula:



96. A benzylated CIPAMFYLLINE™ or analog thereof compound according to claim 94 of the formula:



97. A method for treating or preventing an inflammatory condition or disease in a patient, comprising administering to the patient in need thereof an amount of a benzylated PDE4 inhibitor, where the amount is effective to treat or prevent the inflammatory condition or disease of the patient, and where the benzylated PDE4 inhibitor is prepared by a method according to any one of claims 1-59.

98. A method for treating or preventing an inflammatory condition or disease in a patient, comprising administering to the patient in need thereof an amount of a compound according to any one of claims 60-96, where the amount is effective to treat or prevent the inflammatory condition or disease of the patient.

99. A method of claims 97 or 98 wherein the PDE4 inhibitor inhibits an enzyme selected from phosphodiesterase 4 A, B, C or D or any combination thereof including all splice variants of PDE4 A, B, C and D.

100. A method of claims 97 or 98 wherein the inflammatory condition or disease is an autoimmune condition or disease.

101. A method of claims 97 or 98 wherein the inflammatory condition or disease involves acute or chronic inflammation of bone and/or cartilage compartments of joints.

102. A method of claims 97 or 98 wherein the inflammatory condition or disease is an arthritis selected from rheumatoid arthritis, gouty arthritis or juvenile rheumatoid arthritis.

103. A method of claims 97 or 98 wherein the inflammatory condition or disease is asthma.

104. A method of claims 97 or 98 wherein the condition or disease is associated with the dysregulation of T-cells.

105. A method of claims 97 or 98 wherein the condition or disease is associated with elevated levels of inflammatory cytokines.

106. A method of claim 105 wherein the inflammatory cytokine is IL-2, IL-4 or IL-5.

107. A method of claim 105 wherein the inflammatory cytokine is IFN- γ

108. A method of claim 105 wherein the inflammatory cytokine is TNF- α .

109. A method of claims 97 or 98 wherein the inflammatory condition or disease is multiple sclerosis.

110. A method of claims 97 or 98 wherein the inflammatory condition or disease is pulmonary sarcoidosis.

111. A method of claims 97 or 98 wherein the inflammatory condition or disease is ocular inflammation or allergy.

112. A method of claims 97 or 98 wherein the inflammatory condition or disease is an inflammatory bowel disease.

113. A method of claim 112 wherein the inflammatory bowel disease is Crohn's disease or ulcerative colitis.

114. A method of claims 97 or 98 wherein the inflammatory condition or disease is an inflammatory cutaneous disease.

115. A method of claim 114 wherein the inflammatory cutaneous disease is psoriasis or dermatitis.

116. A method of claims 97 or 98 wherein the inflammatory condition or disease is chronic obstructive pulmonary disease (COPD), bronchitis, emphysema or acute respiratory distress syndrome (ARDS).

117. A method for treating or preventing a disease or condition in a patient, where the disease or condition is associated with pathological conditions that are modulated by inhibiting enzymes associated with secondary cellular messengers, the method comprising administering to the patient in need thereof an amount of a benzylated PDE4 inhibitor, wherein the amount is effective to treat or prevent a disease or condition associated with pathological conditions that are modulated by inhibiting enzymes associated with secondary cellular messengers, and wherein the benzylated PDE4 inhibitor is prepared by a process according to any one of claims 1-59.

118. A method for treating or preventing a disease or condition in a patient, where the disease or condition is associated with pathological conditions that are modulated by inhibiting enzymes associated with secondary cellular messengers, the method comprising administering to the patient in need thereof an amount of a compound according to any one of claims 60-96, wherein the amount is effective to treat or prevent a disease or condition associated with pathological conditions that are modulated by inhibiting enzymes associated with secondary cellular messengers.

119. A method of claims 117 or 118 wherein the enzyme is a cyclic AMP phosphodiesterase.

120. A method of claims 117 or 118 wherein the enzyme is phosphodiesterase 4.

121. A method of treating or preventing transplant rejection in a patient, the method comprising administering to the patient in need thereof an amount of a benzylated PDE4 inhibitor, where the amount is effective to treat or prevent transplant rejection in the patient, and wherein the benzylated PDE4 inhibitor is prepared by a process according to any one of claims 1-59.

122. A method of treating or preventing transplant rejection in a patient, the method comprising administering to the patient in need thereof an amount of a compound according to any one of claims 60-96, where the amount is effective to treat or prevent transplant rejection in the patient.

123. A method of claims 121 or 122 wherein the rejection is due to graft versus host disease.

124. A method of treating or preventing uncontrolled cellular proliferation in a patient, the method comprising administering to the patient in need thereof an amount of a benzylated PDE4 inhibitor, where the amount is effective to treat or prevent uncontrolled cellular proliferation in the patient, and wherein the benzylated PDE4 inhibitor is prepared by a process according to any one of claims 1-59.

125. A method of treating or preventing uncontrolled cellular proliferation in a patient, the method comprising administering to the patient in need thereof an amount of a compound according to any one of claims 60-96, where the amount is effective to treat or prevent uncontrolled cellular proliferation in the patient.

126. A method of claims 124 or 125 wherein the uncontrolled cellular proliferation is caused by a cancer selected from leukemia and solid tumors.

127. A method of treating or preventing conditions associated with the central nervous system (CNS) in a patient, the method comprising administering to the

patient in need thereof an amount of a benzylated PDE4 inhibitor, where the amount is effective to treat or prevent conditions associated with the central nervous system (CNS) in the patient, and wherein the benzylated PDE4 inhibitor is prepared by a process according to any one of claims 1-59.

128. A method of treating or preventing conditions associated with the central nervous system (CNS) in a patient, the method comprising administering to the patient in need thereof an amount of a compound according to any one of claims 60-96, where the amount is effective to treat or prevent conditions associated with the central nervous system (CNS) in the patient.

129. A method of claims 127 or 128 wherein the condition associated with the central nervous system (CNS) is depression.

130. A method of claims 127 or 128 wherein the condition is long-term memory potentiation and learning enhancement.

131. A method of treating or preventing diseases in a patient, the disease associated with viral infection, the method comprising administering to the patient in need thereof an amount of a benzylated PDE4 inhibitor, where the amount is effective to treat or prevent diseases associated with viral infection in the patient, and wherein the benzylated PDE4 inhibitor is prepared by a process according to any one of claims 1-59.

132. A method of treating or preventing diseases in a patient, the disease associated with viral infection, the method comprising administering to the patient in need thereof an amount of a compound according to any one of claims 60-96, where the amount is effective to treat or prevent diseases associated with viral infection in the patient.

133. The method of claims 131 or 132 wherein the viral infection is due to the human immunodeficiency virus (HIV) and the disease is acquired immunodeficiency syndrome (AIDS).

134. A method of treating or preventing diseases in a patient, the disease associated with infection by a parasite, the method comprising administering to the patient in need thereof an amount of a benzylated PDE4 inhibitor, where the amount is effective to treat

or prevent diseases associated with infection of the patient by a parasite, and wherein the benzylated PDE4 inhibitor is prepared by a process according to any one of claims 1-59.

135. A method of treating or preventing diseases in a patient, the disease associated with infection by a parasite, the method comprising administering to the patient in need thereof an amount of a compound according to any one of claims 60-96, where the amount is effective to treat or prevent diseases associated with infection of the patient by a parasite.

136. The method of claims 134 or 135 wherein the parasitic infection is due to the trypanosoma brucei and the disease is African sleeping sickness disease.

137. A method of treating or preventing cystic fibrosis in a patient, the method comprising administering to the patient in need thereof an amount of a benzylated PDE4 inhibitor, where the amount is effective to treat or prevent cystic fibrosis in the patient, where the amount is effective to treat or prevent cystic fibrosis, and wherein the benzylated PDE4 inhibitor is prepared by a process according to any one of claims 1-59.

138. A method of treating or preventing cystic fibrosis in a patient, the method comprising administering to the patient in need thereof an amount of a compound according to any one of claims 60-96, where the amount is effective to treat or prevent cystic fibrosis.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
20 September 2001 (20.09.2001)

PCT

(10) International Publication Number
WO 01/68600 A3

(51) International Patent Classification⁷: **C07D 207/26**,
233/32, 239/10, 473/04, 473/06, 213/75, C07C 271/60,
255/46, A61K 31/4015, 31/4166, 31/513, 31/52, 31/44,
31/325, 31/277

Road, Richmond, British Columbia V6X 1M5 (CA).
SHEN, Yaping [CA/CA]; 21-2458 Pitt River Road, Port
Coquitlam, British Columbia V3C 6J7 (CA).

(21) International Application Number: PCT/CA01/00345

(74) Agents: **NASSIF, Omar, A. et al.**; Gowling Lafleur Hen-
derson LLP, Suite 4900, Commerce Court West, Toronto,
Ontario M5L 1J3 (CA).

(22) International Filing Date: 16 March 2001 (16.03.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/190,337 16 March 2000 (16.03.2000) US

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM,
HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK,
LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,
MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL,
TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(71) Applicant (*for all designated States except US*): **IN-
FLAZYME PHARMACEUTICALS LIMITED**
[CA/CA]; Suite 425, 5600 Parkwood Way, Richmond,
British Columbia V6V 2M2 (CA).

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **LAUENER,**
Ronald, W. [CA/CA]; 818 Massey Street, New Westmin-
ster, British Columbia V3L 4S9 (CA). **BURGOYNE,**
David, L. [CA/CA]; 4715 53A Street, Delta, British
Columbia V4K 3V4 (CA). **REBSTEIN, Patrick, J.**
[CA/CA]; 2659 West 14th Avenue, Vancouver, British
Columbia V6K 2W8 (CA). **MACKENZIE, Lloyd, F.**
[CA/CA]; 10864 129th Street, Surrey, British Columbia
V3T 3N8 (CA). **ZHOU, Yuanlin** [CA/CA]; 11253 Daniels

Published:

— with international search report

(88) Date of publication of the international search report:
28 February 2002

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

WO 01/68600 A3

(54) Title: BENZYLATED PDE4 INHIBITORS

(57) Abstract: The present invention is directed to a method for reducing the emetogenic effects of PDE inhibitors, and more particularly is directed to compounds having PDE4 inhibition activity with little or no emetogenic side-effects, and chemical methods including benzylation for preparing such compounds. A benzyl group may be attached to either a carbon or nitrogen atom of a PDE4 inhibitor. Suitable benzylation chemistry is to extract a hydrogen from a PDE4 inhibitor, preferably with a base, and then react the resulting nucleophilic PDE4 inhibitor with a benzylating agent, e.g. benzyl bromide or a derivative thereof.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/CA 01/00345

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D207/26 C07D233/32 C07D239/10 C07D473/04 C07D473/06
C07D213/75 C07C271/60 C07C255/46 A61K31/4015 A61K31/4166
A61K31/513 A61K31/52 A61K31/44 A61K31/325 A61K31/277

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D C07C A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 95 00139 A (SMITHKLINE BEECHAM CORP ;TORPHY THEODORE J (US); BARNETTE MARY S () 5 January 1995 (1995-01-05) table 1, compound 8 page 6, line 6 -page 7, line 22 page 2, line 13 -page 4, line 3 ---	23,24, 26-31, 63-66, 97,98
X	CHRISTENSEN, S.B. ET AL: "1,4-Cyclohexanecarboxylates: Potent and Selective Inhibitors of Phosphodiesterase 4 for the Treatment of Asthma" JOURNAL OF MEDICINAL CHEMISTRY, vol. 41, 1998, pages 821-835, XP002173934 page 825, column 2, paragraph 3 -page 826, column 1, paragraph 1 table 2 compounds 2-5 --- -/-	23,24, 26-31, 63-66, 97,98

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

3 August 2001

Date of mailing of the international search report

14.11.2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3018

Authorized officer

Seitner, I

INTERNATIONAL SEARCH REPORT

International Application No

PCT/CA 01/00345

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 A61P29/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 94 12461 A (PFIZER ; DUPLANTIER ALLEN J (US); EGGLEER JAMES F (US); MARFAT ANTHO) 9 June 1994 (1994-06-09) example 31 claims 9-12 ---	23,24, 26-31, 63-66, 97,98
X	WO 92 19594 A (SMITHKLINE BEECHAM CORP) 12 November 1992 (1992-11-12) examples 1-58 claims 11,12 --- -/--	23,24, 26-31, 63-66, 97,98

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- "Z" document member of the same patent family

Date of the actual completion of the international search

3 August 2001

Date of mailing of the international search report

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Seitner, I

INTERNATIONAL SEARCH REPORT

International Application No

PCT/CA 01/00345

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	MARIVET M C ET AL: "INHIBITION OF CYCLIC ADENOSINE-3',5'-MONOPHOSPHATE PHOSPHODIESTERASE FROM VASCULAR SMOOTH MUSCLE BY ROLIPRAM ANALOGUES" JOURNAL OF MEDICINAL CHEMISTRY, AMERICAN CHEMICAL SOCIETY, WASHINGTON, US, vol. 32, no. 7, 1989, pages 1450-1457, XP002039410 ISSN: 0022-2623 table 1 page 1452, column 1 ---	23,24, 26-30, 63-65
X	WO 93 07141 A (SMITHKLINE BEECHAM CORP) 15 April 1993 (1993-04-15) page 20 -page 21 examples 7-9,12-15 claims 1,5-9 ---	23,24, 26-31, 63-66, 97,98
P,X	MACDONAL,D. ET AL: "Hunting the Emesis and Efficacy Targets of PDE4 Inhibitors: Identification of the Photoaffinity Probe 9-(3-azidophenyl)-6-[(4-iodo-1H-1-imidazolyl)methyl]quinoline (APIIMQ)" JOURNAL OF MEDICINAL CHEMISTRY, vol. 43, 30 September 2000 (2000-09-30), pages 3820-3823, XP002173935 page 3820, column 2, paragraph 7 -page 3821, column 1, paragraph 1 compound 6 table 1 ---	23,24, 26-31, 63-66, 97,98
X	US 4 950 674 A (YANNI JOHN M ET AL) 21 August 1990 (1990-08-21) preparation 105 ---	63,64
X	BARFACKER L ET AL: "Rhodium(I)-catalysed Hydrocarbonylation and Silylcarbonylation Reactions of Alkynes in the Presence of Primary Amines Leading to 2-Pyrrolidinones and 4-Silylated 1-Aza-1,3-butadienes" TETRAHEDRON, ELSEVIER SCIENCE PUBLISHERS, AMSTERDAM, NL, vol. 54, no. 18, 30 April 1998 (1998-04-30), pages 4493-4506, XP004122050 ISSN: 0040-4020 example 4 --- -/--	63

INTERNATIONAL SEARCH REPORT

In International Application No
PCT/CA 01/00345

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>ROWLEY, M. ET AL: "4-Heterocyclylpiperidines as Selective High-Affinity Ligands at the Human Dopamine D4 Receptor" JOURNAL OF MEDICINAL CHEMISTRY, vol. 40, 1997, pages 2374-2385, XP002173936 example 50</p> <p>---</p>	63
X	<p>KUSTERS E ET AL: "Influence of temperature on the enantioseparation of rolipram and structurally related racemates on Chiracel-OD" JOURNAL OF CHROMATOGRAPHY A, ELSEVIER SCIENCE, NL, vol. 737, no. 2, 21 June 1996 (1996-06-21), pages 333-337, XP004039003 ISSN: 0021-9673 figure 1</p> <p>---</p>	63
X	<p>DATABASE CHEMABS [Online] CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; SUGDEN, J. K.: "Pyrrolidine analog of diphenatil and related substances" retrieved from STN Database accession no. 83:37487 XP002173939 CAS RN: 55470-23-0; 55470-24-1 abstract & PHARM. ACTA HELV. (1974), 49(7-8), 268-9,</p> <p>---</p>	63
X	<p>EP 0 381 235 A (EISAI CO LTD) 8 August 1990 (1990-08-08) page 40, ex (3) page 42, ex (2) page 44, ex (7) page 47, ex (11)</p> <p>---</p>	60,61
A	<p>PIAZ, V.D.: "Novel Heterocyclic-Fused Pyridazinones as Potent and Selective Phosphodiesterase IV Inhibitors" JOURNAL OF MEDICINAL CHEMISTRY, vol. 40, 1997, pages 1417-1421, XP002173937 abstract compounds 10a-d, 11a, 11b page 1418, column 1, paragraph 4 -column 2, paragraph 4</p> <p>---</p> <p style="text-align: center;">-/--</p>	23,97,98

INTERNATIONAL SEARCH REPORT

In. ational Application No
PCT/CA 01/00345

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	KARLSSON J-A ET AL: "PHOSPHODIESTERASE 4 INHIBITORS FOR THE TREATMENT OF ASTHMA" EXPERT OPINION ON THERAPEUTIC PATENTS, ASHLEY PUBLICATIONS, GB, vol. 7, 1997, pages 989-1003, XP000872389 ISSN: 1354-3776 page 994 -page 995 page 998 ---	1,23,97, 98
A	DUPLANTIER A J ET AL: "Biarylcarboxylic acids and -amides: inhibition of phosphodiesterase type IV versus [3H]rolipram binding activity and their relationship to emetic behavior in the ferret" JOURNAL OF MEDICINAL CHEMISTRY, AMERICAN CHEMICAL SOCIETY, WASHINGTON, US, vol. 39, 1996, pages 120-125, XP002150381 ISSN: 0022-2623 abstract compound 18 table 1 ---	1,23,97, 98
A	CRESPO,M.I. ET AL: "Design, Synthesis, and Biological Activities of New Thieno[3,2-d]pyrimidines as Selective Type 4 Phosphodiesterase Inhibitors" JOURNAL OF MEDICINAL CHEMISTRY, vol. 41, 1998, pages 4021-4035, XP002173938 page 4022, column 1, line 1 page 4025 -page 4026 compounds 4,37-39 -----	1,23,97, 98

INTERNATIONAL SEARCH REPORT

International application No.
PCT/CA 01/00345

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Although claims 97-138 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

see further information PCT/ISA/210

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-11 (all partially); 12-15; 23-28 (all partially); 29-31; 60-66; 97-138 (all partially)

Method for improving the therapeutic ratio of PDE4 inhibitors by benzylation of a PDE4 inhibitor according to claims 1 and 23, wherein the PDE4-inhibitor is Rolipram or an analog thereof as well as the compounds according to claims 60 and 63, and their pharmaceutical uses and compositions according to claims 97-138.

2. Claims: 1-3 (all partially); 5-11 (all partially); 16-22; 23-25 (all partially); 27-28 (all partially); 67-72; 97-138 (all partially)

Method for improving the therapeutic ratio of PDE4 inhibitors by benzylation of a PDE4 inhibitor according to claims 1 and 23, wherein the PDE4-inhibitor is Ariflo or an analog thereof as well as the compounds according to claim 67, and their pharmaceutical uses and compositions according to claims 97-138.

3. Claims: 1 (partially); 3-11 (all partially); 23-28 (all partially); 32-35; 73-75; 97-138 (all partially)

Method for improving the therapeutic ratio of PDE4 inhibitors by benzylation of a PDE4 inhibitor according to claims 1 and 23, wherein the PDE4-inhibitor is WAY-PDA-641 or an analog thereof as well as the compounds according to claim 73, and their pharmaceutical uses and compositions according to claims 97-138.

4. Claims: 1 (partially); 3-11 (all partially); 23-28 (all partially); 36-42; 76-81; 97-138 (all partially)

Method for improving the therapeutic ratio of PDE4 inhibitors by benzylation of a PDE4 inhibitor according to claims 1 and 23, wherein the PDE4-inhibitor is Ro-20-1724 or an analog thereof as well as the compounds according to claim 76, and their pharmaceutical uses and compositions according to claims 97-138.

5. Claims: 1 (partially); 3-11 (all partially); 23-28 (all partially); 43-46; 82-84; 97-138 (all partially)

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Method for improving the therapeutic ratio of PDE4 inhibitors by benzylation of a PDE4 inhibitor according to claims 1 and 23, wherein the PDE4-inhibitor is RP 73401 or an analog thereof as well as the compounds according to claim 82, and their pharmaceutical uses and compositions according to claims 97-138.

6. Claims: 1 (partially); 3-11 (all partially); 23-28 (all partially); 47-50; 85-97; 97-138 (all partially)

Method for improving the therapeutic ratio of PDE4 inhibitors by benzylation of a PDE4 inhibitor according to claims 1 and 23, wherein the PDE4-inhibitor is CP-80,633-A or an analog thereof as well as the compounds according to claim 85, and their pharmaceutical uses and compositions according to claims 97-138.

7. Claims: 1 (partially); 3 (partially); 5-11 (all partially); 23-28 (all partially); 51-57; 88-93; 97-138 (all partially)

Method for improving the therapeutic ratio of PDE4 inhibitors by benzylation of a PDE4 inhibitor according to claims 1 and 23, wherein the PDE4-inhibitor is Arofylline or an analog thereof as well as the compounds according to claim 88, and their pharmaceutical uses and compositions according to claims 97-138.

8. Claims: 1 (partially); 3 (partially); 5-11 (all partially); 23-28 (all partially); 58-59; 94-96; 97-138 (all partially)

Method for improving the therapeutic ratio of PDE4 inhibitors by benzylation of a PDE4 inhibitor according to claims 1 and 23, wherein the PDE4-inhibitor is Cipamfylline or an analog thereof as well as the compounds according to claim 94, and their pharmaceutical uses and compositions according to claims 97-138.

9. Claims: 1 (partially); 3-11 (all partially); 23-28 (all partially); 97-138 (all partially)

Method for improving the therapeutic ratio of PDE4 inhibitors by benzylation of a PDE4 inhibitor according to claims 1 and 23, wherein the PDE4 inhibitor is not covered by any of the compounds of the above metioned subjects 1-8

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/CA 01/00345

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9500139 A	05-01-1995	AU 7174794 A EP 0710109 A JP 8511805 T US 6143782 A ZA 9404324 A	17-01-1995 08-05-1996 10-12-1996 07-11-2000 10-05-1995
WO 9412461 A	09-06-1994	AU 673569 B AU 5539694 A BR 9307570 A CA 2150812 A CN 1094028 A CZ 9501417 A EP 0672031 A FI 935379 A HU 65928 A IL 107758 A JP 8501318 T JP 3100984 B NO 952178 A NZ 257955 A PL 309257 A US 5814651 A ZA 9308978 A	14-11-1996 22-06-1994 25-05-1999 09-06-1994 26-10-1994 15-11-1995 20-09-1995 03-06-1994 28-07-1994 20-11-1997 13-02-1996 23-10-2000 01-08-1995 28-05-1996 02-10-1995 29-09-1998 01-06-1995
WO 9219594 A	12-11-1992	AU 1917092 A CA 2102106 A CN 1067244 A EP 0584208 A IE 921376 A JP 6507405 T MX 9202090 A PT 100441 A ZA 9203210 A	21-12-1992 03-11-1992 23-12-1992 02-03-1994 04-11-1992 25-08-1994 01-11-1992 30-09-1993 31-03-1993
WO 9307141 A	15-04-1993	AU 2869092 A MX 9205875 A PT 100947 A ZA 9207787 A	03-05-1993 01-07-1993 28-02-1994 20-07-1993
US 4950674 A	21-08-1990	US 4810713 A AU 629534 B AU 2982289 A NZ 228114 A ZA 8901224 A AU 580444 B AU 6247286 A DK 243086 A EP 0235463 A EP 0228893 A GR 862023 A IE 59726 B JP 62148468 A NZ 217773 A PH 23981 A ZA 8604458 A	07-03-1989 08-10-1992 24-08-1989 29-01-1991 31-10-1990 12-01-1989 25-06-1987 21-06-1987 09-09-1987 15-07-1987 24-12-1986 23-03-1994 02-07-1987 26-02-1990 09-02-1990 25-02-1987
EP 0381235 A	08-08-1990	JP 2928288 B JP 3118361 A	03-08-1999 20-05-1991

INTERNATIONAL SEARCH REPORT

Information on patent family members

ii. National Application No

PCT/CA 01/00345

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0381235 A		AT 92043 T	15-08-1993
		AU 619943 B	06-02-1992
		AU 4901390 A	01-11-1990
		CA 2009162 A	03-08-1990
		CN 1044652 A	15-08-1990
		DE 69002338 D	02-09-1993
		DE 69002338 T	16-12-1993
		DK 381235 T	04-10-1993
		ES 2058620 T	01-11-1994
		HU 53867 A	28-12-1990
		JP 2288855 A	28-11-1990
		JP 2928307 B	03-08-1999
		KR 9203892 B	18-05-1992
		NO 173988 C	02-03-1994
		NZ 232334 A	25-06-1991
		PT 93045 A	31-08-1990
		RU 2024506 C	15-12-1994
		US 5444083 A	22-08-1995
